

Organ regeneration is no longer a distant dream

July 12 2018




Figure 1: The left-right asymmetric rotation and cell chirality in the *Drosophila* embryonic hindgut. Credit: Osaka University

Many organs arise from simple sheets and tubes of cells. During development, these sheets bend and deform into the more complex final shape of the organ. This can be seen, for example, in the hindgut of fruit flies (*Drosophila*), which is an organ equivalent to our intestines.

The *Drosophila* embryonic hindgut first forms as a bilaterally symmetric structure and then rotates 90° counterclockwise as viewed from the posterior to become asymmetric. Before the hindgut [rotation](#), cells in the hindgut tilt leftward and are asymmetric. This structure cannot be superimposed on its mirror image, so this asymmetric [structure](#) is called cell chirality.

As cell chirality disappears after the rotation, it was thought that cell chirality was related to this hindgut rotation. Previous studies, however, were performed using fixed samples, so the cellular dynamic mechanism by which cellular chirality was converted into an LR asymmetric organ remained unknown.

Researchers led by Kenji Matsuno at Osaka University used live imaging of the *Drosophila* embryonic hindgut and computer simulations to clarify that a novel cellular behavior called "cell sliding" was important for the LR asymmetric morphogenesis of the organ. Their research results were published in *eLife*.

Lead Author Mikiko Inaki said, "We found that when the hindgut rotated, the cells changed their position relative to the [cells](#) located below them (posterior in the embryo) by sliding leftward. We named this movement 'cell sliding.' In an inversion mutant showing inverted cell [chirality](#) and hindgut rotation, cell sliding occurred in the opposite direction to that found in the wild-type. Thus, we confirmed that cell sliding was important for the LR asymmetric rotation of the hindgut."

This group's achievements will deepen an understanding of the

mechanism behind the morphogenesis of organs that have tubular structures, such as the digestive tracts and heart. It is also anticipated that their research results will be applied to the regeneration of organs in the future.




Figure 2: Cell sliding revealed by live imaging and computer simulations. Credit:

Osaka University

More information: Mikiko Inaki et al. Chiral cell sliding drives left-right asymmetric organ twisting, *eLife* (2018). [DOI: 10.7554/eLife.32506](https://doi.org/10.7554/eLife.32506)

Provided by Osaka University

Citation: Organ regeneration is no longer a distant dream (2018, July 12) retrieved 2 May 2024 from <https://phys.org/news/2018-07-regeneration-longer-distant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.