


# Rice plants evolve to adapt to flooding

July 12 2018



Temporal morphological changes of rice in deepwater conditions. Credit: Takeshi Kuroha, Keisuke Nagai, and Motoyuki Ashikari


Although water is essential for plant growth, excessive amounts can

waterlog and kill a plant. In South and Southeast Asia, where periodic flooding occurs during the rainy season, the water depth can reach several meters for many months.

Rice varieties known as "deepwater [rice](#)" have developed a unique strategy to ensure their own survival. Deepwater rice grows normally in shallow water but in heavy floods increases its height in keeping with rising water levels, to enable the [plants](#) to ride out lengthy floods.

A research team comprising Takeshi Kuroha at Tohoku University, Motoyuki Ashikari at Nagoya University, Susan R. McCouch at Cornell University and colleagues in Japan and the U.S.A., have discovered a gene in rice that is critical to its survival in flood conditions. They have also shed light on its molecular function and evolutionary history.

The research group identified the SD1 (SEMIDWARF1), as a key gene responsible for the deepwater rice's response. The SD1 encodes a biosynthesis enzyme of gibberellin—a plant hormone. The gene orchestrates the deepwater rice response via a unique gain-of-function allele. When submerged, rice accumulate ethylene, a gaseous plant hormone. Deepwater rice amplify a signaling relay in which the SD1 gene is transcriptionally activated by an ethylene-responsive transcription factor, OsEIL1a.




Molecular mechanism in deepwater rice. Credit: Takeshi Kuroha, Keisuke Nagai, and Motoyuki Ashikari

The resulting SD1 protein directs increased synthesis of gibberellins, largely one of gibberellin species, GA<sub>4</sub>, which promote vertical growth in the plant. Further analysis revealed that this conditionally functional variation evolved first in a wild ancestor and was then a target of selection during the domestication of cultivated rice adapted to deepwater environments in Bangladesh.

The SD1 gene is well-known as the Green Revolution gene in rice, where a loss-of-function allele of SD1 confers short plant height, providing lodging resistance and increases the harvest index, generating

greater grain yields under high input agricultural systems (Figure 3- left).

A transcriptional gain-of-function allele of the same gene enables deepwater rice to adapt to flooding via the opposite phenotypic response—an increase in plant height. The ability of SD1 to function in such diverse roles in cultivated rice highlights the inherent plasticity of plant response to its environment.



The SD1 gene enables the plant to adapt to different environments. Credit: Takeshi Kuroha, Keisuke Nagai, and Motoyuki Ashikari

"Extreme weather events caused by climate change could affect food production worldwide," said Kuroha. "Farmers will need to diversify their methods and the cryptic genetic variation found in wild rice [genes](#) may offer adaptive solutions for growing resilient crops."

**More information:** [DOI: 10.1126/science.aat1577](https://doi.org/10.1126/science.aat1577) T. Kuroha et al., "Ethylene-gibberellin signaling underlies adaptation of rice to periodic flooding," *Science* (2018). [science.sciencemag.org/cgi/doi ... 1126/science.aat1577](https://science.sciencemag.org/cgi/doi/10.1126/science.aat1577)

Provided by Tohoku University

Citation: Rice plants evolve to adapt to flooding (2018, July 12) retrieved 9 April 2024 from <https://phys.org/news/2018-07-rice-evolve.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---