

# LED lights reduce seabird death toll from fishing by 85 percent, research shows

July 10 2018

---



Guanay cormorant stuck in a net. Credit: Andrew F Johnson

Illuminating fishing nets with low-cost lights could reduce the terrible impact they have on seabirds and marine-dwellers by more than 85 per cent, new research has shown.

A team of international researchers, led by Dr. Jeffrey Mangel from the University of Exeter, has shown the number of birds caught in gillnets can be drastically reduced by attaching green battery-powered light-emitting diodes (LEDs).

For the study, the researchers compared 114 pairs of gillnets—which are anchored in fixed positions at sea and designed to snare fish by the gills—in fishing waters off the coast of Peru.

They discovered that the nets fitted with the LEDs caught 85 per cent fewer guanay cormorants—a native diving bird that commonly becomes entangled in nets—compared with those without lights.

Coupled with previous research conducted by the same team, that showed LED lighting also reduced the number of [sea turtles](#) caught in fishing nets by 64 per cent, the researchers believe the lights offer a cheap, reliable and durable way to dramatically reduce the capture and death of birds and [turtles](#), without reducing the intended catch of fish.

The research is published in the journal *Royal Society Open Science* on Wednesday, July 11 2018.

Lead author Dr. Mangel, from the Centre for Ecology and Conservation at the University's Penryn Campus, said: "We are very encouraged by the results from this study.



Fishing net with LED lights. Credit: ProDelphinus

"It shows us that we may be able to find cost-effective ways to reduce bycatch of multiple taxa of protected species, and do so while still making it possible for fishers to earn a livelihood."

Peru's gillnet fleet comprises the largest component of the nation's small-scale fleet and is conservatively estimated to set 100,000km of net per year in which thousands of turtles and seabirds will die as "bycatch" or unintentionally.

The innovative study, carried out in Sechura Bay in northern Peru, saw the LED lights attached at regular intervals to commercial fishing gillnets which are anchored to the bottom of the water. The nets are left in situ from late afternoon until sunlight, when the fishermen collect their haul.

The researchers used 114 pairs of nets, each typically around 500-metres in length. In each pair, one was illuminated with light-emitting diodes (LEDs) placed every ten metres along the gillnet floatline. The other net in the pair was the control and not illuminated.

The control nets caught 39 cormorants, while the illuminated nets caught just six.

A previous study, using the same LED technology, showed they also reduced the number of sea turtles also caught in gillnets. Multiple populations of sea turtle species use Peruvian coastal waters as foraging grounds including green, olive ridley, hawksbill, loggerhead and leatherback.

For that study, the researchers found that the control nets caught 125 green turtles while illuminated nets caught 62. The target catch of guitarfish was unaffected by the net illumination. They are now working with larger fisheries in Peru and with different coloured lights to see if the results can be repeated and applied with more [critically endangered species](#).

Professor Brendan Godley, who is an author of the study and Marine Strategy Lead for the University of Exeter, said: "It is satisfying to see the work coming from our Exeter Marine Ph.D.s leading to such positive impact in the world. We need to find ways for coastal peoples to fish with the least impact on the rest of the biodiversity in their seas."



**More information:** Illuminating gillnets to save seabirds and the potential for multi-taxa bycatch mitigation, *Royal Society Open Science*, [rsos.royalsocietypublishing.org ... /10.1098/rsos.180254](https://rsos.royalsocietypublishing.org/.../10.1098/rsos.180254)

Provided by University of Exeter

Citation: LED lights reduce seabird death toll from fishing by 85 percent, research shows (2018, July 10) retrieved 18 June 2024 from <https://phys.org/news/2018-07-seabird-death-toll-fishing-percent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.