


# Surprising similarity between stripy black holes and high-temperature superconductors

July 23 2018, by Erik Arends

---



Credit: Leiden Institute of Physics

Scientists don't understand how some materials become superconducting at relatively high temperatures. Leiden physicists have now found a surprising connection with auxiliary black holes. It enables researchers to apply knowledge of black holes to the mystery of high-temperature superconductivity. The new study is published in *Nature Physics*.

Superconductivity, discovered in Leiden in 1911, is exploited for many modern applications, such as MRI scanners and particle accelerators. These are based on the phenomenon whereby electric currents flow without resistance at temperatures close to absolute zero. Unfortunately, continuous cooling requires lots of energy, so physicists all over the world are looking for a way to make superconducting materials that operate at higher temperatures.

All relatively high-temperature superconductors now known are based on so-called Mott insulators. These form when electrons are stuck in their crystal lattice nodes, exactly one per node. They turn into superconductors after extra electrons are injected. Researchers don't understand why this happens on a fundamental level. Knowing this could enable even higher-temperature superconductors that are cheaper to keep sufficiently cool.

As the superconductor is formed, the disparity between the number of electrons and the number of available nodes within the crystal lattice causes a stripy pattern, much like the moving Moiré patterns seen on TV when an old-fashioned computer screen is filmed. But why? This is a key question in understanding Mott insulators.

Physicists are looking for the answer in an unexpected direction—they hypothesize that the dodgy electrons in high-temperature [superconductors](#) behave in some ways similarly to auxiliary [black holes](#). Leiden physicists Alexander Krikun, Koenraad Schalm and Jan Zaanen together with Tomas Andrade from the University of Barcelona have now found the same stripy pattern in a similar discrepancy between auxiliary 'wavy' black holes and a [crystal lattice](#). This confirms the hypothesis and means that knowledge about black holes can apply to better understanding [high-temperature superconductivity](#).

**More information:** Tomas Andrade, Alexander Krikun , Koenraad Schalm & Jan Zaanen, 'Doping the holographic Mott insulator', *Nature Physics*. [arxiv.org/abs/1710.05791](https://arxiv.org/abs/1710.05791)

Provided by Leiden Institute of Physics

Citation: Surprising similarity between stripy black holes and high-temperature superconductors

(2018, July 23) retrieved 10 April 2024 from <https://phys.org/news/2018-07-similarity-stripy-black-holes-high-temperature.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.