

Tropical treetops are warming, putting sensitive species at risk

July 26 2018



Changing temperatures in the forest canopies of Panama's Barro Colorado Island could mean major consequences for overall forest health. Credit: Stephanie Pau

The vaulted canopies that tower above Earth's rich tropical forests could be especially vulnerable to climate change-related temperature increases,


according to a new report from Florida State University researchers.

In a study published in the journal *Ecosphere*, scientists found temperatures in the [forest](#) canopies of Panama's Barro Colorado Island exceeded maximum [air temperatures](#) by as much as 7 degrees Celsius. Their results suggest that as air temperatures continue to climb, canopy temperatures could increase at a 40 percent higher rate—a shift of potentially major consequence for tropical forest health.

"Tropical forests are among the most productive places on Earth—they account for about 30 percent of terrestrial productivity," said study author Stephanie Pau, an assistant professor in the Department of Geography. "These forests are vulnerable to climate change and climbing temperatures."

Tropical species like those in the canopies of Barro Colorado Island evolved to survive within a narrow range of equatorial temperatures, and are therefore hypothesized by scientists to be especially sensitive to minute temperature change.

In this study, researchers analyzed the effect of canopy temperature increase on the forests' gross primary productivity, or the amount of carbon the forests absorb and process through photosynthesis. They found productivity rates increase with warming temperatures but begin to slow at temperatures between 28 and 29 degrees Celsius (82.4 and 84.2 degrees Fahrenheit), and to decline at [high temperatures](#) exceeding 31 degrees Celsius (87.8 degrees Fahrenheit).



Using a refined thermal imaging system, Pau was able to record detailed breakdowns of canopy temperature changes over several months. Credit: Stephanie Pau

"We wanted to determine the consequences of rising canopy temperatures for the ecological functioning of these forests," Pau said. "We show that as temperatures warm, there might be a reduction in productivity."

Using a refined thermal imaging system, Pau and her team captured detailed snapshots of the canopies' thermal profile every five minutes for several months. This continuous, precise monitoring showed that temperatures were highly variable—within half a degree of the mean temperature less than 20 percent of the time—indicating significant and potentially disruptive fluctuations.

The high spatial resolution imaging also allowed researchers to discriminate between different forest surfaces. Evergreen canopies, for example, cooled faster than deciduous canopies after reaching peak temperatures, and tree bark reached warmer temperatures than flower

clusters.

This classified breakdown of forest surfaces revealed a dynamic variation in temperature change. Overall temperature increase, accompanied by this pronounced variability, suggests that some canopy-dwelling species may be at particular risk.



Perched in towers high above the treetops, Pau and her team had a panoramic view of the forest canopies. Credit: Stephanie Pau

"There are so many different organisms that use and partition the canopies in different ways," Pau said. "For example, ant species in

tropical forests are extremely sensitive to bark temperatures. The diversity of canopy dwelling species will be affected by canopy temperatures—and it's not just how hot it will get, but it's also the temperature variability within a canopy."

Rising air temperatures are routinely cited by scientists and lay people alike as a sobering effect of our rapidly changing climate, but Pau's research demonstrates that these straightforward measurements may not tell the whole story. If the atmosphere continues to warm, she and her team discovered, [canopy](#) temperatures will warm at a rate of 1.4 degrees Celsius for every 1-degree increase in air [temperature](#).

For plant and animal species perfectly suited for survival within a narrow and stable range of temperatures, this kind of dramatic increase could cause considerable harm.

"When people talk about climate change, we usually hear about air temperatures, but we should keep in mind that tropical forest canopies are even more vulnerable than air temperatures might indicate," Pau said.

Pau's study is titled "Tropical Forest Temperature Thresholds for Gross Primary Productivity."

More information: Stephanie Pau et al, Tropical forest temperature thresholds for gross primary productivity, *Ecosphere* (2018). [DOI: 10.1002/ecs2.2311](https://doi.org/10.1002/ecs2.2311)

Provided by Florida State University

Citation: Tropical treetops are warming, putting sensitive species at risk (2018, July 26) retrieved

5 April 2024 from <https://phys.org/news/2018-07-tropical-treetops-sensitive-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.