

Researchers describe unique molecular activity in water-ethanol mixtures

July 2 2018



Droplet of ionic liquid (Red) with active holes (white holes) in a water-ethanol solvent (Blue). The holes propel themselves inside the IL droplet. Credit: Rei Kurita



Researchers from Tokyo Metropolitan University have observed the formation of holes that move by themselves in droplets of ionic liquids (IL) sitting inside water-ethanol mixtures. This complex phenomenon is driven by an interplay between how ionic liquids dissolve, and how the boundary around the droplet fluctuates. Self-driven motion is a key feature of active matter, materials that use ambient energy to self-propel, with potential applications in drug delivery and nano-machine propulsion.

It is well known that substances like <u>water</u> and ethanol mix very well at room temperature; how well alcoholic beverages mix depends on the environment in which the mixture takes place, including such factors as temperature and pressure.

However, dissolution can take a complex turn. A team led by Associate Professor Rei Kurita, Department of Physics, Tokyo Metropolitan University, was studying how an ionic liquid (IL) dissolved in a mixture of water and ethanol. Ionic liquids are liquids composed entirely of ions in ambient conditions; properties like their resistance to drying and ability to dissolve otherwise difficult materials make them a promising solvent, with possible applications in industrial processes such as battery production, pharmaceuticals and recycling.

The team placed a small droplet of IL at the bottom of a mixture of ethanol and water. With the temperature and particular ratio of ethanol to water they used, they expected a boundary or interface to form between the IL and the water-ethanol above it, and for the two to mix gradually. Yet, what they saw was startling: Over time, holes emerged inside the IL droplet, and the holes could propel themselves inside the droplet.



They found that this curious phenomenon was the result of how the composition of the water-ethanol mixture naturally fluctuated around the interface. The conditions were such that the mixture is close to a critical point at which small variations in composition can have major consequences. In this case, they were enough to promote local mixing of the IL into the water-ethanol mixture; the unique way in which ILs interact with water resulted in further local changes in composition, leading to a positive feedback loop, or instability. The effect was so drastic that they produced variations in the surface tension, causing the surface to become spontaneously bumpy, form holes, and generate the large flows required to move them around. These holes have been dubbed active holes.

Their discovery paves the way for a broad new class of synthetic active matter, materials that can spontaneously take energy from the surroundings and convert it into motion. With possible applications in <u>drug delivery</u> and propulsion at the nanometer scale, this new phenomenon might inspire investigations into novel industrial uses as well as further accelerate academic interest in active phenomena.

More information: Noriko Oikawa et al, Active hole generation in a liquid droplet dissolving into a binary solvent, *Soft Matter* (2018). DOI: 10.1039/c8sm00357b

Provided by Tokyo Metropolitan University

Citation: Researchers describe unique molecular activity in water-ethanol mixtures (2018, July 2) retrieved 2 May 2024 from https://phys.org/news/2018-07-unique-molecular-water-ethanol-mixtures.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.