

Abandoned farmlands enrich bird communities

August 3 2018



Abandoned wet farmland in central Hokkaido in Japan. Credit: Forestry and Forest Products Research Institute and Hokkaido University

Agriculture and conversion of pristine lands into urban or industrial areas have exerted immense pressure on natural biota due to habitat destruction and fragmentation in industrialized countries around the world. But since the 1900s, farmlands have been increasingly abandoned



due to the decline in domestic agriculture and, in some countries, a decline in population. This yields an opportunity for abandoned farmlands to be used as rehabilitation zones for grassland, wetland, and forest animals. However, it has so far remained unclear how valuable for sustaining specific animal communities farmlands, abandoned farmlands, and natural habitats are relative to each other.

Hokkaido University's Futoshi Nakamura and his collaborators including Yuichi Yamaura of Forestry and Forest Products Research Institute, published in the journal *Agriculture, Ecosystems and Environment,* studied <u>bird species</u> distribution over an area in central Hokkaido, Japan. The group then correlated abundance and <u>species richness</u> of different bird communities with different degrees of farmland abandonment and the landscape structure. Mapped back onto the landscape of the known composition, this allowed them to evaluate the potential to conserve different bird communities if existing farmlands in a given area were to be abandoned.

The team came to the conclusions that abandoned farmlands, especially when not surrounded by forests, benefit wetland and grassland birds more than other bird communities. However, how well a farmland can be reclaimed depends centrally on the intensity of farmland usage before abandonment. Furthermore, different landscape structure properties affect bird communities in differential ways. For example, a large proportion of newly abandoned farmland within a given area can increase species richness of wetland bird species, favoring rare bird species; on the other hand, farmland abandonment in open areas can increase abundance of wetland species, favoring common species.

The effects of landscape structure on animal communities are complex and, as a result, different studies report conflicting results. However, this recent study clearly indicates that abandoned farmlands can be used for the preservation of wetland and grassland fauna. In addition, it untangles



some of the factors underlying complicated interactions, thus yielding valuable insights for choosing land plots for specific conservation efforts.



Abandoned dry farmland in central Hokkaido in Japan. Credit: Forestry and Forest Products Research Institute and Hokkaido University



(a) Wetland species



(b) Grassland species

Nakamura's study produced a concrete prediction where different bird groups would be most abundant in the current agricultural landscape. Credit: Hanioka M. et al., Agriculture, Ecosystems and Environment, June 26, 2018

More information: Masashi Hanioka et al, Assessing the landscapedependent restoration potential of abandoned farmland using a hierarchical model of bird communities, *Agriculture, Ecosystems & Environment* (2018). DOI: 10.1016/j.agee.2018.06.014

Provided by Hokkaido University



Citation: Abandoned farmlands enrich bird communities (2018, August 3) retrieved 16 June 2024 from <u>https://phys.org/news/2018-08-abandoned-farmlands-enrich-bird.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.