

OK computer: How AI could help forecast quake aftershocks

August 29 2018



Seismologists have methods to forecast when aftershocks will hit and how strong they will be, but there is more uncertainty about how to predict where they will strike

Lightning might not strike twice, but earthquakes can. And forecasting where aftershocks will hit might now be a little easier thanks to an assist

from artificial intelligence.

Aftershocks can be more destructive than the quakes they follow, making it all the more important for experts to be able to predict them.

But while seismologists have methods to forecast when aftershocks will hit and how strong they will be, there is more uncertainty about how to predict where they will strike.

Hoping to address that, a group of researchers trained a "deep learning" programme with data about tens of thousands of earthquakes and aftershocks to see if they improve predictions.

"The previous baseline for aftershock forecasting has a precision of around three percent across the testing data set. Our neural network approach has a precision of around six percent," said Phoebe DeVries, co-author of the study published in the journal *Nature* on Thursday.

"This approach is more accurate because it was developed without a strongly held prior belief about where aftershocks ought to occur," DeVries, a post-doctoral fellow at Harvard, told AFP.

The researchers used a type of artificial intelligence known as [deep learning](#), which is loosely modelled on the way the human brain makes connections.

The programme allowed the researchers to map relationships "between the characteristics of a large earthquake—the shape of the fault, how much did it slip, and how did it stress the earth—and where aftershocks occurred," said Brendan Meade, professor of earth and planetary science at Harvard, and a study co-author.

The researchers tested the network by holding back a quarter of their

data set, and feeding the remaining information into the programme.

They then tested how well the programme predicted the aftershock locations of the 25 percent of cases it hadn't been fed.

They found six percent of the areas the programme identified as high-risk did in fact experience aftershocks, up from three percent using existing methods.

Analysing the research, Gregory Beroza, a professor of geophysics at Stanford University, cautioned it "might be premature to infer... an improved physical understanding of [aftershock](#) triggering".

In an article published in *Nature* alongside the study, he said the research had focused on only one set of changes caused by earthquakes that can affect where aftershocks occur.

"Another reason for caution is that the authors' analysis relies on factors that are fraught with uncertainty," Beroza wrote.

DeVries acknowledged that additional factors affect where aftershocks occur and that there is "much more to be done".

"We definitely agree that this work is a motivating beginning, rather than an ending," she said.

And Beroza said the research had established a "beachhead" for additional study into how [artificial intelligence](#) could help forecasting.

"The application of machine-learning methods has the potential to extract meaning from these large and complex sources of information, but we are still in the early stages of this process."

More information: Phoebe M. R. DeVries et al. Deep learning of aftershock patterns following large earthquakes, *Nature* (2018). [DOI: 10.1038/s41586-018-0438-y](https://doi.org/10.1038/s41586-018-0438-y)

© 2018 AFP

Citation: OK computer: How AI could help forecast quake aftershocks (2018, August 29) retrieved 26 April 2024 from <https://phys.org/news/2018-08-ai-quake-aftershocks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.