


# Algal blooms a threat to small lakes and ponds, too

August 23 2018, by Misti Crane

---



Credit: CC0 Public Domain

Harmful algae isn't just a problem for high-profile bodies of water—it

poses serious, toxic threats in small ponds and lakes as well, new research has found.

A team of researchers from The Ohio State University examined [water](#) samples from two dozen ponds and small lakes in rural Ohio and found plenty of cause for concern, with particularly high levels of toxins at one lake.

Toxins from algae can cause skin rashes, intestinal problems and damage to the liver and nervous system. Fertilizers common to agriculture—including nitrogen and phosphorous—create an environment in which [harmful algae](#) can flourish.

The researchers said that the way farmers manage runoff could play a significant role in creating water bodies that are ripe for harmful [algal blooms](#). A primary concern is [tile drainage](#), a widely used agricultural approach to removing excess water from the soil below the surface. That water—and the nutrients found in it—are rerouted, often toward ponds on farm property, said study co-author Seungjun Lee, a postdoctoral researcher in [environmental health sciences](#) at Ohio State.

"A lot of people and government agencies are paying attention to larger lakes, including Lake Erie, but these smaller bodies of water are also used for recreation, fishing and irrigation," he said.

The study was published recently in the journal *Environmental Science & Technology*.

The research team, led by Ohio State's Jiyoung Lee, analyzed samples from the 24 bodies of water over a three-month period in late summer 2015.

Ten of the sites had detectable levels of microcystins, toxins produced

by freshwater cyanobacteria during algal blooms.

One site had repeated instances of microcystin concentrations above recreational guidelines set by the Ohio Environmental Protection Agency, and so the research team paid particular attention to the samples from that site.

"Samples from this lake in early July were particularly concerning, as they contained four times the recommended amount of microcystin for recreational use and more than 800 times the recommended level for drinking," Seungjun Lee said.

A [pond](#) or lake with high toxin levels presents a risk to people, pets, farm animals, wildlife (including fish) and crops and could benefit from routine monitoring and work to lower the risk of algal blooms, he said. The researchers did not name the [lake](#) in question, because it is privately owned.

Jiyoung Lee said the impact of tile drainage may be elevated in small lakes and ponds, compared to larger lakes.

"Highly concentrated nutrients are being introduced into a smaller volume of water, making small lakes and ponds more sensitive to this influx of phosphorous, nitrogen and other nutrients," she said.

Nitrate and phosphorous are linked to the primary type of toxic microcystin found in the water. Judicious use of fertilizers could help control the algal blooms, as could measures to reduce animal waste contamination of ponds and lakes, Seungjun Lee said.

Though the study concentrated on Ohio agricultural areas, its findings likely apply to many areas throughout the U.S. and the world where agriculture and small lakes and ponds coexist, the researchers said.

Provided by The Ohio State University

Citation: Algal blooms a threat to small lakes and ponds, too (2018, August 23) retrieved 27 April 2024 from <https://phys.org/news/2018-08-algal-blooms-threat-small-lakes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.