

Ecosystems are getting greener in the Arctic

August 21 2018, by Theresa Duque



Credit: Wikimedia Commons

In recent decades, scientists have noted a surge in Arctic plant growth as a symptom of climate change. But without observations showing exactly when and where vegetation has bloomed as the world's coldest areas warm, it's difficult to predict how vegetation will respond to future warming. Now, researchers at the U.S. Department of Energy's



Lawrence Berkeley National Laboratory (Berkeley Lab) and UC Berkeley have developed a new approach that may paint a more accurate picture of Arctic vegetation and our climate's recent past – and future.

In a study published online Aug. 20 in *Nature Climate Change*, the researchers used satellite images taken over the past 30 years to track – down to a pixel representing approximately 25 square miles – the ebb and flow of <u>plant growth</u> in cold areas of the northern hemisphere, such as Alaska, the Arctic region of Canada, and the Tibetan Plateau.

The 30-year historic satellite data used in the study were collected by the National Oceanic and Atmospheric Administration's Advanced Very High Resolution Radiometer. The data was processed by Boston University, and is hosted on NEX – the NASA Earth Exchange data archive.

At first, the satellite data showed what they expected – that as Arctic climates warmed, tree and plant growth increased. After comparing these observations with state-of-the-art <u>climate</u> models developed for CMIP5 – the Coupled Model Intercomparison Project Phase 5 – what they discovered next surprised them.

Their data analysis revealed that 16 percent of Earth's vegetated land where plant growth was limited by cold temperatures three decades ago is no longer predominantly temperature-limited today, a result that was not reproduced by the CMIP5 models tested. "Our findings suggest that CMIP5's predictions may have significantly underestimated changes in the Arctic ecosystem, and climate models will need to be improved to better understand and predict the future of the Arctic," said first author Trevor Keenan, a faculty scientist in Berkeley Lab's Earth & Environmental Sciences Area and an assistant professor in UC Berkeley's department of Environmental Science, Policy, and Management.





Current and predicted changes in areas around the world where vegetation responds strongly to warming. Shaded areas represent areas where vegetation growth was limited by cold temperatures at the start of the observational record (1982-1986) but was not by the corresponding year. Representative Concentration Pathway (RCP) 4.5 is a carbon emissions target to limit temperature increases to 2 degrees Celsius, while RCP 8.5 represents a scenario where carbon emissions are not limited and temperature increases exceed 2 degrees Celsius. Credit: Trevor Keenan and William Riley/Berkeley Lab

Keenan and Riley used the <u>satellite data</u> to build a new observational benchmark that quantifies the growing expanse of vegetated land in the northern hemisphere. They also estimated changes in the proportion of the Earth's surface where plant growth will no longer be limited by cold temperatures over the 21st century. Keenan and Riley project that by the year 2100, only 20 percent of vegetated land in the <u>northern hemisphere</u> will still be limited by cold conditions that have been in place there for



centuries; the remaining 80 percent will no longer experience sufficiently <u>cold temperatures</u>, and with earlier springs, plants will grow sooner, in unexpected places and to an unexpected degree.

"Although the greening might sound like good news as it means more carbon uptake and biomass production, it represents a major disruption to the delicate balance in cold ecosystems," said Keenan. "Temperatures will warm sufficiently so that new species of trees could move in and compete with vegetation that had previously dominated the landscape. This change in vegetation would also affect insects and animals that relied on native vegetation for food."

Scientists collaborating through the World Climate Research Programme developed the CMIP5 models to help researchers around the world gain a better understanding of the relationship between carbon emissions and global warming, among other goals. International consortiums such as the IPCC (International Panel on Climate Change) have also used CMIP5 projections to inform policy decisions. Keenan said that while the CMIP5 models provided researchers with a broad overview of the problem, they do not always accurately represent the important roles plants play in reflecting light back into the atmosphere, sending water back into the atmosphere, and absorbing carbon dioxide.

"No one has looked at high-latitude systems from this angle before as they are very complex, but they're important as they control multiple feedbacks to the Earth system," said co-author William Riley, a senior scientist in Berkeley Lab's Earth & Environmental Sciences Area.

Now that Keenan and Riley have established a standard approach for assessing <u>climate models</u>, they plan to explore how they can use more advanced statistical techniques, such as machine learning, to quantify how soil organic matter properties, atmospheric carbon dioxide, wildland fires, and temperature, will affect climate in the 21st century.



More information: T. F. Keenan et al. Greening of the land surface in the world's cold regions consistent with recent warming, *Nature Climate Change* (2018). DOI: 10.1038/s41558-018-0258-y

Provided by Lawrence Berkeley National Laboratory

Citation: Ecosystems are getting greener in the Arctic (2018, August 21) retrieved 3 May 2024 from <u>https://phys.org/news/2018-08-ecosystems-greener-arctic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.