

Extreme weather in Europe linked to less sea ice and warming in the Barents Sea

August 30 2018



Credit: AI-generated image ([disclaimer](#))

Yueng-Djern Lenn, *Bangor University*; Benjamin Barton, *Bangor University*, and Camille Lique, *Institut Français de Recherche pour l'Exploitation de la Mer (Ifremer)*

The cold, remote Arctic Ocean and its surrounding marginal seas have

experienced climate change at a rate not seen at lower latitudes. Warming air, land and sea temperatures, and large declines in seasonal Arctic sea ice cover are all symptoms of the changing Arctic climate. Although these changes are occurring in relatively remote locations, there is growing evidence to link Arctic sea ice retreat to increasingly erratic weather patterns over the northern hemisphere.

As sea ice declines, areas of open water increase, allowing the ocean to lose more heat to the atmosphere. Heat lost from the ocean to the atmosphere reduces the atmospheric pressure which provides more energy to storms and increases their cloud content through evaporation.

Water flowing north from the Atlantic Ocean provides a major source of heat to the Arctic Ocean and surrounding continental shelf seas. While the Atlantic Water (the particular water mass in the Arctic ocean) carries enough heat to melt all the floating Arctic sea ice in less than five years, it is currently insulated from the surface by a lighter, fresher layer of water over most of the central Arctic Ocean.

However, this paradigm appears to be changing. North of Svalbard, Atlantic Water heat has been mixed up towards the surface, resulting in increased surface heat lost to the atmosphere over the ever greater area of open ocean. This change has recently been shown to enhance the rate of sea ice loss eastwards.



Location of the Barents Sea. Credit: Wikimedia, CC BY-SA


Barents Sea changes

A key Arctic region for Atlantic Water heat exchange with the atmosphere is the Barents Sea. Atlantic Water flowing east through the Barents Sea Opening – between Bear Island, and northern Norway – remains exposed to the atmosphere as it circulates through the central

Barents Sea. It gradually cools and becomes fresher (due to sea ice melting) as it moves eastwards to the Kara Sea.

In the Barents Sea, sea ice forms every autumn and melts in late spring/summer. In the northern part of the sea, a north-south change from cold to warm sea surface temperatures signals the presence of the Polar Front, which separates cold Arctic water from the warm Atlantic water. The meeting of the two water masses, its location and the temperature difference across it reflects changes in Barents Sea circulation.

During years with low seasonal sea ice concentrations (when there's more heat loss from more exposed open water), the north-south differences in atmospheric temperatures across the Barents Sea are reduced. These conditions have been linked to [wintertime cyclones](#) travelling further south into western Europe, instead of their tendency to move eastwards towards Siberia, as well as more frequent [cold winter extremes at middle latitudes](#).



Winter-averaged sea surface temperature and sea ice extent as observed in the Barents Sea by satellites from 2005 and 2016. Credit: Author provided

Ice and weather

[For our recent study](#), we looked at satellite measurements of sea ice and sea surface temperature, to determine how ocean and ice conditions have evolved between 1985 and the end of 2016. We found that prior to 2005, sea ice extended south of the Polar Front every winter, but that since 2005 this has not been the case.

At the same time, the [sea surface temperature](#) difference across the Polar Front has increased, with southern temperatures increasing at a faster rate than those to the north. The average between 1985 and 2004

was -1.2°C in the north and 1.5°C in the south, while between 2005 and 2016 it was -0.6°C in the north and 2.6°C in the south. Clearly, from 2005 the Barents Sea has become too warm for sea ice to exist south of the Polar Front. The question then is why is the Barents Sea getting warmer?

[Long-term oceanographic measurements](#) of water temperature and salinity near the Barents Sea Opening have shown that inflowing Atlantic Water temperatures have increased over the last 30 years, with what appears to be a [small but persistent rise around 2005](#) – likely to be due to upstream changes in the North Atlantic sources (though it must be noted that our study did not explore this question). An impact of the warmer water entering the Barents Sea is a warmer atmosphere, which in turn insulates the warmer surface water allowing the Atlantic Water [heat](#) to penetrate further to the north, preventing winter sea ice formation and import (that is sea ice that has formed farther north that has drifted southwards) to the region south of the Polar Front.

We believe that this represents a long-term shift in the climate of the Barents Sea, a region already identified as influential on lower-latitude European weather. Furthermore, we believe that the 2005 regime shift we observed over the Barents Sea may have contributed to the increasingly frequent extreme weather events experienced over Europe in the past decade or so.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Extreme weather in Europe linked to less sea ice and warming in the Barents Sea (2018, August 30) retrieved 24 April 2024 from <https://phys.org/news/2018-08-extreme-weather-europe->

[linked-sea.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.