

The fading ghost of a long-dead star

August 2 2018, by Calla Cofield



Thin, red veins of energized gas mark the location of the supernova remnant HBH3 in this image from NASA's Spitzer Space Telescope. The puffy, white feature in the image is a portion of the star forming regions W3, W4 and W5. Infrared wavelengths of 3.6 microns have been mapped to blue, and 4.5 microns to red. The white color of the star-forming region is a combination of both wavelengths, while the HBH3 filaments radiate only at the longer 4.5 micron wavelength. Credit: NASA/JPL-Caltech/IPAC

Thin, red veins of energized gas mark the location of one of the larger supernova remnants in the Milky Way galaxy in this image from NASA's Spitzer Space Telescope.

A supernova "remnant" refers to the collective, leftover signs of an exploded star, or supernova. The red filaments in this image belong to a supernova remnant known as HBH 3 that was first observed in 1966 using radio telescopes. Traces of the remnant also radiate optical light. The branches of glowing material are most likely molecular gas that was pummeled by a shockwave generated by the supernova. The energy from the explosion energized the molecules and caused them to radiate infrared light.

The white, cloud-like formation also visible in the image is part of a complex of [star-forming regions](#), simply named W3, W4 and W5. However, those regions extend far beyond the edge of this image. Both the white star-forming regions and the red filaments are approximately 6,400 light years away and lie inside our Milky Way galaxy.

HBH 3 is about 150 light-years in diameter, ranking it amongst the largest known supernova remnants. It is also possibly one of the oldest: Astronomers estimate the original explosion may have happened anywhere from 80,000 to one million years ago.

In 2016, NASA's Fermi Gamma-Ray Telescope detected very high-energy light—called gamma rays—coming from the region near HBH 3. This emission may be coming from gas in one of the neighboring star-forming regions, excited by powerful particles emitted by the [supernova](#) blast.

The Spitzer Space Telescope is one of NASA's four Great

Observatories—along with the Hubble Space Telescope, the Chandra X-ray Observatory and the Compton Gamma-Ray Observatory—and will celebrate its 15th birthday in space on Aug. 25. Spitzer sees the universe in [infrared light](#), which is slightly less energetic than the optical light we can see with our eyes. In this image, taken in March 2010, infrared wavelengths of 3.6 microns have been mapped to blue, and 4.5 microns to red. The white color of the star-forming region is a combination of both wavelengths, while the HBH3 filaments radiate only at the longer 4.5-micron wavelength.

Provided by Jet Propulsion Laboratory

Citation: The fading ghost of a long-dead star (2018, August 2) retrieved 9 April 2024 from <https://phys.org/news/2018-08-ghost-long-dead-star.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--