


# Can microswimmers swim through gel?

August 28 2018



Artist's illustration of a three-sphere microswimmer in a polymer gel network.  
Credit: Shigeyuki Komura

Researchers from Tokyo Metropolitan University have studied how microswimmers, like bacteria or sperm, swim through fluids with both

solid and liquid-like properties, e.g., gels. They found that subtle changes in a swimmer's features, its structure, and how it moves invoke a dramatically different response from the fluid. They also discovered that the similarity in size between the structure of the fluid and the swimmer led to a wide range of interesting behavior.

Swimming is a tricky business for the microorganism. A dip in the pool might not seem that difficult, but at microscopic scales, or at low Reynolds numbers, the effect of [fluid](#) viscosity imposes severe constraints on swimming. Yet, nature achieves it; microswimmers play vital roles in a wide range of phenomena, including sperm motility and the active motion of bacteria.

To understand swimmers, previous studies have focused on minimal models of [swimmer](#) behavior in uniform fluids. A particularly popular model is the so-called "three-sphere microswimmer," a string of three microscopic spheres attached to each other by arms; the string can be propelled by pumping the arms backward and forward in a liquid. This simple sequence overcomes the limitations of the "scallop" theorem of Purcell, which says that motion that looks the same when played backwards ([time-reversal symmetry](#)), like a scallop opening and closing, cannot be used for locomotion.

But what about the fluid? In the case of sperm traveling through cervical mucus to reach eggs in mammalian reproduction, the mucus is an example of soft matter, where the internal structure, in this case made of sugars and proteins, responds in a complex fashion to the motion of the swimmer. To address this issue, a team consisting of Kento Yasuda and Associate Professor Shigeyuki Komura of Tokyo Metropolitan University and Ryuichi Okamoto, a lecturer at Okayama University, studied how three-sphere microswimmers behave in a structured fluid, a polymer gel e.g. jelly.

Their analysis revealed that there were broadly two mechanisms for achieving motion, one by breaking time-reversal symmetry, the other by modulating the amplitudes in the beating of the two arms of the swimmer. With the latter, it was found that swimming could be achieved without breaking the former symmetry, a loophole in the scallop theorem. Through further detailed analysis, they succeeded in deriving expressions for how the velocity of the swimmer was related to how a structured fluid resists the motion of a swimmer. Interestingly, they found that when swimmers were larger than the mesh size of the gel, there was greater resistance with faster beating, a somewhat counterintuitive conclusion.

This work marks significant progress in bringing a popular minimal swimmer model closer to experimentally relevant cases, including the beating of hairs ("cilia") on cells and the motility of bacteria. It may also see application to more exotic scenarios, e.g. the locomotion of robots through debris after landslides. The study has been published online in the journal *Europhysics Letters*.

**More information:** Kento Yasuda et al. A three-sphere microswimmer in a structured fluid, *EPL (Europhysics Letters)* (2018). [DOI: 10.1209/0295-5075/123/34002](https://doi.org/10.1209/0295-5075/123/34002)

Provided by Tokyo Metropolitan University

Citation: Can microswimmers swim through gel? (2018, August 28) retrieved 1 May 2024 from <https://phys.org/news/2018-08-microswimmers-gel.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.