


To stop mosquitoes, target their young

August 9 2018, by Ayleen Barbel Fattal



East-central Miami showing the 5 km² Wynwood area where both naled and the larvicide Bti were applied. Red markers indicate locations where mosquito traps were placed in that area. Credit: Florida International University

Bacterial larvicide, not standard insecticides, are the best solution for mosquito control, according to new research by Florida International University biologist Philip Stoddard.

When it comes to the *Aedes aegypti* mosquito, which is responsible for Zika transmission, controlling populations at the larval stage is particularly effective. Overall, larvicides are more effective and a safer long-term solution as opposed to insecticides that target adult populations, Stoddard said.

"Old-style control using adulticides is not the silver bullet we need for controlling today's urban [mosquitoes](#), and maybe not even part of the arsenal," Stoddard said. "We need to develop and adopt new biological control methods before the next big disease hits us."

Results from the recent study show larvicides were responsible for steady declines in the *Aedes aegypti* populations, with some populations dropping as much as 90 percent. In those cases, naturally occurring bacteria found in soil was used that only targets mosquito larvae.

The study actually analyzed three methods of mosquito control deployed in the summer of 2016 when the first case of locally transmitted Zika was reported in South Florida. Miami-Dade County Mosquito Control used a mix of insecticides targeting adults and larvicides to combat the mosquitoes. Population data was determined based on mosquito counts

from traps placed in the Zika zones of Wynwood and Miami Beach.



Miami Beach neighborhoods of South Beach where Zika appeared first and Mid-Beach into which Zika expanded. Red markers indicate trap locations. Credit: Florida International University

Results showed insecticides targeting adult mosquitoes had no significant effect on [population](#) counts in Miami Beach. Naled, a potent and controversial [insecticide](#), was among the adulticides used to combat the Zika-spreading mosquitoes. Derived from a class of chemicals developed as nerve agents during WWII, naled has raised concerns about its potential harm to non-targets including people and pollinating insects. After three days of spraying in Wynwood, the *Aedes aegypti* populations were virtually identical to pre-spray levels, according to Stoddard. Average counts were actually 7 percent higher after those first three days.

Stoddard says fundamental changes need to occur in how communities manage mosquito control.

He recommends:

- Urban mosquito control should happen before a disease strikes
- More research is needed to better understand how urban mosquitoes adapt
- Scientists and statisticians should be involved in control efforts from the start
- Urban mosquito control should employ more larvicides and not rely on conventional adulticides

These findings were published this month in *Acta Tropica*.

More information: Philip K. Stoddard. Managing *Aedes aegypti* populations in the first Zika transmission zones in the continental United States, *Acta Tropica* (2018). [DOI: 10.1016/j.actatropica.2018.07.031](https://doi.org/10.1016/j.actatropica.2018.07.031)

Provided by Florida International University

Citation: To stop mosquitoes, target their young (2018, August 9) retrieved 1 May 2024 from <https://phys.org/news/2018-08-mosquitoes-young.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.