


How plant-rotting bacteria steal iron to survive

August 2 2018



Mechanism for the import of the plant protein ferredoxin (red) across the bacterial outer membrane and into the periplasm of *Pectobacterium*. Credit: Trevor Lithgow, [10.1371/journal.pbio.2006026](https://doi.org/10.1371/journal.pbio.2006026)

In a new study, researchers identify important new insights into a survival mechanism of the bacteria that cause rotting in certain plants, including some highly invasive weeds. The study, publishing on August 2 in the Open-Access journal *PLOS Biology*, demonstrates for the first time exactly how the bacterium *Pectobacterium* obtains the iron vital to its survival and replication: by pirating it from iron-bearing proteins in

the host plants. The study was led by Dr. Rhys Grinter and Prof. Trevor Lithgow at the Biomedicine Discovery Institute of Monash University. The team used comparative genomics to forensically track the origins of an enzyme, called FusC, that turned out to be the key factor for the import and acquisition of iron.

The authors propose that plant ferredoxin, a [protein](#) loaded with iron, is imported into the bacterial cell through a membrane channel called FusA; when it arrives in the cell interior FusC grabs it and dismembers the ferredoxin to release the iron. "This is the first example of bacterium taking a protein from its host, importing it into the bacterial cell then processing it inside the cell," Dr. Grinter said. *Pectobacterium* is of increasing scientific interest for its potential as a [biological control agent](#) against [invasive weeds](#) such as *Allium triquetrum* (angled onion), which can smother native ground flora such as orchids, lilies and grasses, leaving large infested areas bare of natural vegetation.

"Knowing exactly how the bacterium obtains iron adds to the knowledge needed to maximise the potential of *Pectobacterium* as a biocontrol agent," Dr. Grinter said. The study also reveals new insights into the evolution of this bacteria, finding that it 're-evolved' the iron-importing mechanism in a way that paralleled the protein import pathways evolved in mitochondria and chloroplasts, the energy centers of plant cells.

"Rhys' work is a remarkable example of how fundamental discoveries in biological science have far-reaching impact," senior author Prof. Trevor Lithgow said. "The study not only elucidates the mechanism of how this bacterium imports [iron](#)-loaded proteins, with its obvious practical applications, but also reveals a new paradigm for how bacteria evolve protein transport pathways in general. This will include the bacteria that infect humans and animals, not just [plants](#)," he said.

More information: Rhys Grinter et al, FusC, a member of the M16

protease family acquired by bacteria for iron piracy against plants, *PLOS Biology* (2018). [DOI: 10.1371/journal.pbio.2006026](https://doi.org/10.1371/journal.pbio.2006026)

Provided by Public Library of Science

Citation: How plant-rotting bacteria steal iron to survive (2018, August 2) retrieved 3 May 2024 from <https://phys.org/news/2018-08-plant-rotting-bacteria-iron-survive.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--