

Research reveals effective method to control algae growth on Hawaiian coral reefs

August 8 2018



Divers using the Super Sucker to manually remove *Eucheuma clade* (algae; left) and outplanting juvenile *Tripneustes gratilla* (urchins; right). Credit: DLNR/DAR

Researchers with the State of Hawai'i Division of Aquatic Resources (DAR) and the Hawai'i Institute of Marine Biology (HIMB) at the University of Hawai'i at Mānoa found a management approach that



combining manual removal and outplanting native urchin was effective at reducing invasive, reef smothering macroalgae by 85% on a coral reef off O'ahu, Hawai'i.

Globally, the health of coral reefs is threatened due to rising ocean temperatures and ocean acidification. However, local factors such as invasive macroalgae also pose a serious risk to coral reefs—monopolizing <u>reef</u> habitats, and overgrowing and smothering native species, such as corals.

Brian J. Neilson, Acting Administrator at DAR, and Chris Wall, doctoral candidate at HIMB in the UH Mānoa School of Ocean and Earth Science and Technology (SOEST), and others tested a novel approach to curbing the abundance of invasive macroalgae on the coral reefs of Kāne'ohe Bay, O'ahu. First, divers manually removed invasive macroalgae with the assistance of an underwater vacuum system, "The Super Sucker." Then, hatchery-raised juvenile <u>sea urchins</u> (the Hawaiian native collector urchin, Tripneustes gratilla) were outplanted to graze on invasive algae to control regrowth.

In total, the team removed over 40,000 pounds of invasive macroalgae, outplanted 99,000 sea urchins, and treated nearly six acres of reef area over two years. During this period, invasive macroalgae declined in response to treatments, and importantly, there were no observed negative effects to important reef calcifiers such as corals and crustose coralline algae.





Outplanted adult *T. gratilla* surrounded by *Gracilaria salicornia* and *Acanthophora spicifera* (left) and adult *T. gratilla* surrounded by *Eucheuma clade* (right). Credit: Brian Neilson

Unfortunately, there are often limited options for reducing invasive macroalgae without causing further environmental damage. Prior to this study, scientists at DAR, UH Mānoa and the Nature Conservancy showed the manual removal/urchin herbivory method worked at reducing invasive macroalgae in the laboratory and in small enclosures on the reef.

"This management approach is the first of its kind at the reef-scale," said Wall. "Our research shows promise as an effective mean to reduce invasive macroalgae with minimal environmental impact, while also incorporating a native herbivore to regulate a noxious invasive species."

"Coral reefs are an important part of the economy, culture, sustenance and recreation of Hawai'i," said Neilson. "Local action is instrumental in supporting the resilience of coral reefs. This study provides an important tool that can assist in the management and conservation of coral reefs."





Top row: Before (left) and immediately after (right) manual removal of *Eucheuma clade* revealing crustose coralline algae. Bottom row: Before (left) and immediately after (right) removal of *Eucheuma clade* revealing live and dead coral. Credit: Brian Neilson

"The surprise was just how effective this approach was at reducing invasive macroalgae over the two-year period," said Wall. "We were able to successfully leverage the rigorous, detailed science of prior studies to assist in scaling the management plan from an aquarium to an entire reef. One of the lessons here is that a well-designed management plan can reap significant benefits and lead to the most effective path forward, both logistically and financially."



DAR is continuing to monitor the reefs of Kāne'ohe Bay and the longterm effects of macroalgae removal and urchin herbivory on <u>coral reefs</u>. In addition, DAR and UH scientists are actively studying the influence of local weather and global climate phenomena, such as the 2014-2016 El Niño and global bleaching episodes, as drivers of changing coral and invasive <u>macroalgae</u> abundance through time.

More information: Brian J. Neilson et al, Herbivore biocontrol and manual removal successfully reduce invasive macroalgae on coral reefs, *PeerJ* (2018). DOI: 10.7717/peerj.5332

Provided by University of Hawaii at Manoa

Citation: Research reveals effective method to control algae growth on Hawaiian coral reefs (2018, August 8) retrieved 17 May 2024 from <u>https://phys.org/news/2018-08-reveals-effective-method-algae-growth.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.