

Rh ensemble catalyst for effective automobile exhaust treatment

August 29 2018



Professor Hyunjoo Lee and PhD candidate Hojin Jeong. Credit: The Korea Advanced Institute of Science and Technology (KAIST)

A KAIST research team has developed a fully dispersed Rh ensemble catalyst (ENS) that shows better performance than commercial diesel oxidation catalyst (DOC). This newly developed ENSs could improve low-temperature automobile exhaust treatment.



Precious metals have been used for various heterogeneous reactions, but it is crucial to maximize efficiency of catalysts due to their high cost. Single-atom catalysts (SACs) have received much attention because it is possible for all of the metal atoms to be used for reactions, yet they do not show catalytic activity for reactions that require ensemble sites.

Meanwhile, hydrocarbons, such as propylene (C3H6) and propane (C3H8) are typical automobile exhaust gas pollutants and must be converted to carbon dioxide (CO2) and water (H2O) before they are released as exhaust. Since the <u>hydrocarbon</u> oxidation reaction proceeds only during carbon-carbon (C-C) or carbon-hydrogen (C-H) bond cleavage, it is essential to secure the metal ensemble site for the catalytic reaction. Therefore, precious metal catalysts with high dispersion and ensemble sites are greatly needed.

To solve this issue, Professor Hyunjoo Lee from the Department of Chemical and Biomolecular Engineering and Professor Jeong Woo Han from POSTECH developed an Rh ensemble catalyst with 100 percent dispersion, and applied it to automobile after-treatment. Having a 100 percent dispersion means that every metal atom is used for the reaction since it is exposed on the surface.





Figure 1. Concept of Rh ensemble catalyst for automobile exhaust treatment. Credit: The Korea Advanced Institute of Science and Technology (KAIST)

SACs also have 100 percent dispersion, but the difference is that ENSs have the unique advantage of having an ensemble site with two or more atoms.

As a result of the experiment, the ENSs showed excellent catalytic performance in CO, NO, propylene, and propane oxidation at low temperatures. This complements the disadvantage of nanoparticle



<u>catalyst</u> (NPs) that perform catalysis poorly at low temperatures due to low metal dispersion, or SACs without hydrocarbon oxidation.

In particular, the ENSs have superior low-temperature activity even better than commercial DOC, hence they are expected to be applied to automobile exhaust treatment.



Figure 2. Structure and performance comparison of single-atom catalyst and ensemble catalyst. Credit: The Korea Advanced Institute of Science and Technology (KAIST)

Professor Lee said, "I believe that the ENSs have given academic contribution for proposing a new concept of metal catalysts, differentiating from conventional SACs and NPs. At the same time, they are of great value in the industry of exhaust treatment catalysts."

This research, led by Ph.D. candidate Hojin Jeong, was published in the *Journal of the American Chemical Society* on July 5.





Figure 3. Energy-dispersive X-ray spectroscopy (EDS) mapping images for SAC, ENS, and NP, respectively (green, Eh; red, Ce). Credit: The Korea Advanced Institute of Science and Technology (KAIST)

More information: Hojin Jeong et al. Fully Dispersed Rh Ensemble Catalyst To Enhance Low-Temperature Activity, *Journal of the American Chemical Society* (2018). DOI: 10.1021/jacs.8b04613

Provided by The Korea Advanced Institute of Science and Technology (KAIST)

Citation: Rh ensemble catalyst for effective automobile exhaust treatment (2018, August 29) retrieved 30 April 2024 from https://phys.org/news/2018-08-rh-ensemble-catalyst-effective-automobile.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.