

Study questions the role of lead poisoning in Franklin Expedition deaths

August 24 2018



In the summer of 1845, under the command of Sir John Franklin, 128 officers and crew aboard the ships HMS Erebus, above, and HMS Terror entered the waters of Arctic North America with the goal of completing the discovery of the Northwest Passage. Credit: National Maritime Museum

A team of investigators from across Canada, including a trio of Western

researchers, have raised serious doubt about the popular belief that lead poisoning played a role in the death of members of the famed Franklin Expedition. The study, Franklin expedition lead exposure: New insights from high resolution confocal X-ray fluorescence imaging of skeletal microstructure, was published today in *PLOS ONE*.

In the summer of 1845, under the command of Sir John Franklin, 128 officers and crew aboard the ships HMS Erebus and HMS Terror entered the waters of Arctic North America with the goal of completing the discovery of the Northwest Passage. Franklin and his crew spent the first winter at Beechey Island, where three crewmen died and were buried.

The following year the ships became stranded in ice off King William Island where they remained until April 1848. By this time the crew, now reduced to 105 men, made a desperate attempt to reach the mainland. Sadly, not one individual survived.

Previous analyses of bone, hair, and soft tissue samples from the remains of crew members found that tissues contained elevated lead levels, suggesting that [lead poisoning](#) may have been a major contribution to their demise.

However, questions remained regarding the timing and degree of exposure to lead and, ultimately, the extent to which the crew members may have been impacted. To address this historical question, the research team investigated three hypotheses to test the theory that lead poisoning was not the primary cause of the crew's deaths:

- If elevated [lead exposure](#) was experienced by the crew during the expedition, the team hypothesized that those sailors who survived longer (King William Island vs. Beechey Island) would exhibit more extensively distributed lead in their bones;

- The team hypothesized that lead levels would be elevated in bone microstructural features forming at or near the time of death, compared with older tissue in the body; and
- Finally, if lead exposure played a significant role in the failure of the expedition the team hypothesized that bone samples would exhibit evidence of higher or more sustained levels of lead than that of a contemporary British naval 19th-Century naval population from Antigua.

Synchrotron-based high resolution confocal X-ray fluorescence imaging in partnership with scientists at the Canadian Light Source synchrotron at the University of Saskatchewan and the Advanced Photon Source was used to visualize lead distribution within bone and dental structures at the micro scale.

The data did not support the first hypothesis as lead distribution within the samples from the two different Franklin sites was similar. Evidence of lead within skeletal microstructural features formed near the time of death lent support to the team's second hypothesis but consistent evidence of a marked elevation in [lead levels](#) was lacking.

Finally, the comparative analysis with the Antigua samples did not support the hypothesis that the Franklin sailors were exposed to an unusually high level of lead compared to navy personnel from the same time period.

Taken all together, the team's skeletal microstructural results do not support the conclusion that lead played a pivotal role in the loss of Franklin and his crew.

More information: Treena Swanston et al. Franklin expedition lead exposure: New insights from high resolution confocal x-ray fluorescence imaging of skeletal microstructure, *PLOS ONE* (2018). [DOI:](#)

[10.1371/journal.pone.0202983](https://doi.org/10.1371/journal.pone.0202983)

Provided by University of Western Ontario

Citation: Study questions the role of lead poisoning in Franklin Expedition deaths (2018, August 24) retrieved 19 April 2024 from <https://phys.org/news/2018-08-role-poisoning-franklin-deaths.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.