


Satellite measurements of the Earth's magnetosphere promise better space weather forecasts

August 10 2018




Bursts of electromagnetic waves named "chorus" captured by the Arase/PWE in the inner magnetosphere. Credit: Kanazawa University

Earth is constantly being hammered by charged particles emitted by the sun with enough power to make life on Earth almost impossible. Life is only possible because Earth's magnetic field traps and deflects these particles, preventing the vast majority of them from ever reaching the planet's surface. The trapped particles bounce back and forth between the North and South poles in complex, ever-changing patterns that are also influenced by intricate and shifting electric fields. When the Van Allen radiation belts in which they travel dip into the atmosphere near the poles, they create the Northern (and Southern) lights. Bursts of these particles can also damage satellites and sensitive equipment on the ground.

It is therefore vital to understand the intricacies of the [radiation belts](#). NASA has launched twin satellites to study the Van Allen belts—however, their orbits only allow them to explore the equatorial regions. This limits our ability to understand flow of particles and prevents us from predicting their effects on all satellites.

To explore regions further from the equator, the Institute of Space and Astronautical Science, a division of the Japan Aerospace Exploration Agency, launched the Arase satellite in 2016. A Japan-based research team centered at Kanazawa University equipped the Arase satellite with multiple sensors for the Plasma Wave Experiment, which probed the electric field and plasma waves in the Earth's inner magnetosphere. Now, they have collected their first set of data from the sensors, which they recently published in *Earth, Planets and Space*.

The Arase consists primarily of electric and magnetic [field](#) detectors covering a wide frequency range; it can also measure plasma/particles in a wide energy range. To improve efficiency, an on-board computer studies the correlations between the fields and the particles before sending only the most important information back to Earth.




Conceptual illustration of wave-particle interaction occurring in the Earth's inner magnetosphere to be explored by the ERG satellite. Credit: (C)JAXA ERG science team

"The Plasma Wave Experiment equipment has passed initial checks and has successfully acquired high-quality data. A huge amount of burst waveform data has been taken, and we should soon know a lot more about mechanisms of wave-particle interactions occurring in the inner magnetosphere than before. Another strength of our project is that we can also compare the [satellite](#) data with data collected simultaneously on the ground. We expect those comparisons will greatly broaden our understanding of this area of science," says first author Yoshiya

Kasahara.

Understanding how electrons and other [particles](#) are hurled out of the magnetosphere toward Earth could be key to predicting such bursts and protecting against them.



Overview of the Arase satellite and configuration of the sensors implemented on the Arase satellite. Credit: (C)JAXA ERG science team

More information: Yoshiya Kasahara et al, The Plasma Wave Experiment (PWE) on board the Arase (ERG) satellite, *Earth, Planets and Space* (2018). [DOI: 10.1186/s40623-018-0842-4](https://doi.org/10.1186/s40623-018-0842-4)

Provided by Kanazawa University

Citation: Satellite measurements of the Earth's magnetosphere promise better space weather forecasts (2018, August 10) retrieved 20 April 2024 from <https://phys.org/news/2018-08-satellite-earth-magnetosphere-space-weather.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.