

# Scientists find coarse resolution models underestimate future Mei-yu rainfall

August 17 2018



Too strong or too weak western North Pacific Subtropical High leads to less Mei-yu rainfall, which is more prone to happen in future projection with a coarse

resolution model. Credit: CHEN Xiaolong

Climate models are indispensable tools for future climate projection, including predictions of the Mei-yu front rainfall, which is the major rain-bearing system of the East Asian summer monsoon. Adaptation to climate change relies heavily on the data of climate projection from the Coupled Model Intercomparison Project (CMIP). However, resolutions of CMIP models such as CMIP5 are generally around 100 to 200 kilometers, which are not high enough to resolve some regional rainfall systems, for instance, the Mei-yu rain belt along the Yangtze River valley. Uncertainty in regional rainfall projection due to model resolution needs to be quantified.

Recently, Dr. Chen Xiaolong and Prof. Zhou Tianjun from the Institute of Atmospheric Physics, Chinese Academy of Sciences, cooperating with Drs. Peili Wu and Malcolm Roberts from Hadley Centre, U.K. Met Office, investigated the effect of model resolution on the Mei-yu [rainfall](#) projection using the Hadley Centre's latest climate model, HadGEM3-GC2. The high-resolution HadGEM3-GC2 (N216, ~60km) projects large increases of summer rainfall under two representative concentration pathway scenarios (RCP8.5 and RCP4.5) whereas the low-resolution (N96, ~130km) shows a decrease. A larger increase of projected Mei-yu rainfall in higher-resolution models is also observed across the CMIP5 ensemble.

Why do climate models employing different resolutions show such large differences in the projected rainfall changes? "Based on moist static energy and moisture budget analysis, we find that the deficient energy moist static energy advection and moisture convergence by stationary eddies (mainly in the meridional direction) in N96 (~130km) is attributed to the projected weaker upward motion and less mei-yu

rainfall," Dr. Chen said.

A large-scale manifestation of the anomalous stationary eddies is the contrasting response to the same warming scenario by the western North Pacific subtropical high (WNPSH), which keeps unchanged in N216 (~60km) but retreats evidently eastward in N96 (~130km) as CMIP5 models. Interactions between high-latitude wave activity and low-latitude convection are dynamical reasons for the distinct responses of WNPSH in high and low resolution HadGEM3-GC2. With unchanged circulation in N216, more moisture in warmer climate will increase the projected Mei-yu rainfall, whereas eastward retreat too much in N96 impedes moist static energy and moisture being transported into the Mei-yu region.

"This investigation highlights the need of high-resolution models in future [climate change](#) projections. While [climate models](#) used in the past CMIPs are generally low resolution models, I am happy to see that in the ongoing CMIP6 there is a high resolution models inter-comparison project, the results will be helpful to further clarifying the resolution-dependence of rainfall projection," said corresponding author Prof. Zhou Tianjun.

**More information:** Xiaolong Chen et al, Potential Underestimation of Future Mei-Yu Rainfall with Coarse-Resolution Climate Models, *Journal of Climate* (2018). [DOI: 10.1175/JCLI-D-17-0741.1](https://doi.org/10.1175/JCLI-D-17-0741.1)

Provided by Chinese Academy of Sciences

Citation: Scientists find coarse resolution models underestimate future Mei-yu rainfall (2018, August 17) retrieved 25 April 2024 from <https://phys.org/news/2018-08-scientists-coarse-resolution-underestimate-future.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.