


Discovering trailing components of a coronal mass ejection

August 27 2018



In the image above, Mercury is seen as the bright dot on the right; the MESSENGER spacecraft was orbiting it. Radio signals from MESSENGER probed the near-solar environment observed. Credit: Tom and Elizabeth Kuiper, (JPL/STAR Prep Academy)

Using Green Bank Observatory in West Virginia, PSI Associate Research Scientist Elizabeth A. Jensen's team observed radio signals from the MESSENGER spacecraft and discovered that solar eruptions known to cause communication disruptions and electrical grid failures as they hit Earth have secondary trailing impacts. Jensen is lead author of "Plasma Interactions with the Space Environment in the Acceleration Region: Indications of CME-trailing Reconnection Regions" that appears in *The Astrophysical Journal*. Co-authors include PSI Senior Scientists Deborah Domingue Lorin and Faith Vilas.

Coronal mass ejections, or CMEs, are powerful, massive [solar eruptions](#) that can trigger geomagnetic storms, Jensen said. CMEs are often associated with solar flares.

"The most powerful CMEs may travel at 2,000 kilometers per second, passing the Earth in seconds. We have discovered reconnection regions trailing behind a CME traveling 750 kilometers per second," Jensen said. "This suggests that the impact of a CME on the Earth's space weather consists of the initial shock from the CME and also secondary shocks from powerful electrical currents and accelerated plasmas trailing in the CME's wake."

These events can affect human activity and technology not only in space, but also within the atmosphere and ultimately on the ground, Jensen said. In 1989, a powerful CME caused problems in satellites orbiting the Earth, the space shuttle Discovery, and electric grids resulting with 6

million people in Canada losing power.

"These secondary regions trailing the CME front have never been seen before because we have had few opportunities to probe this area with spacecraft. The MESSENGER spacecraft orbiting Mercury on the other side of the Sun was in the right place at the right time to observe a CME safely by passing [radio signals](#) through it to the Earth," said Jensen.

"Understanding the three-dimensional structure of these eruptions that strike the Earth is essential to properly preparing for their potential effects."

More information: Elizabeth A. Jensen et al. Plasma Interactions with the Space Environment in the Acceleration Region: Indications of CME-trailing Reconnection Regions, *The Astrophysical Journal* (2018). [DOI: 10.3847/1538-4357/aac5dd](#)

Provided by Planetary Science Institute

Citation: Discovering trailing components of a coronal mass ejection (2018, August 27) retrieved 30 April 2024 from

<https://phys.org/news/2018-08-trailing-components-coronal-mass-ejection.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--