

Towards untangling the 'antennal grabbing' phenomenon in mating cuckoo bees

August 27 2018



Mating in the species *Nomada lathburiana* in the lab, showing the male (top) to spirally entangle the female's antennae with its own. Credit: Matthias Schindler

One can seldom spot a cuckoo bee, whose peculiar kleptoparasitic behaviour includes laying eggs in the nests of a certain host bee species, let alone a couple mating.

Nevertheless, German scientists—Dr. Matthias Schindler, University of Bonn, Michaela Hofmann and Dr. Susanne S. Renner of the University of Munich, and Dr. Dieter Wittmann, recently managed to record copulation in three different [cuckoo bee species](#) in the genus *Nomada*.

Intriguingly, in field and lab settings alike, the observed couples demonstrated the phenomenon the researchers called "antennal grabbing" where the male cuckoo bee winds his antennae around the female's during copulation, thus transferring pheromones. Even though such behaviour is relatively common in Hymenoptera, this is the first known record for the genus *Nomada*.

While the particular biological reason for the "antennal grabbing" in different species remains unsettled, the scientists discuss the phenomenon in view of both previous hypotheses and their own observations in a new paper published in the open access *Journal of Hymenoptera Research*.

The courtship in *Nomada* [cuckoo](#) bee starts with the 'swarming' of males at willow shrubs and gooseberry or their patrolling in groups with males of the *Andrena* or *Melitta* species that will "foster" their offspring. There is no aggression among the males. They were observed to rub their bellies and heads against the grass, in order to leave sexual pheromones, thus marking the "dating spot" for potential mates.



Two males of the species *Nomada flavoguttata* patrolling at a blossom of a common dandelion. Credit: Michaela Hofmann

Earlier chemical studies of *Nomada* bees noted that the mandibular glands of males produce chemical compounds identical with those of their *Andrena* or *Melitta* hosts, leading to the suggestion that the males transfer pheromones that help the females mimic the odor of the host bee, and later enter its nest unnoticed to lay its eggs. An alternative explanation for the "antennal grabbing" is that [males](#) are spraying a substance onto the females to make them unattractive to other potential mates.



Insertion phase of copulation in a couple of the species *Nomada flavoguttata*. Note the male's antennae spirally entangling the female's. Credit: Michaela Hofmann

More information: Matthias Schindler et al, Courtship behaviour in the genus *Nomada* – antennal grabbing and possible transfer of male secretions, *Journal of Hymenoptera Research* (2018). [DOI: 10.3897/jhr.65.24947](https://doi.org/10.3897/jhr.65.24947)

Provided by Pensoft Publishers

Citation: Towards untangling the 'antennal grabbing' phenomenon in mating cuckoo bees (2018, August 27) retrieved 18 May 2024 from <https://phys.org/news/2018-08-untangling-antennal-phenomenon-cuckoo-bees.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.