

# Getting warmer—understanding threats to ocean health

August 16 2018, by Roz Cummins

---



Brad Linsley (center), a paleoclimate scientist at Lamont-Doherty Earth Observatory, and his research team in Panama, March 2018. Credit: Brad Linsley

The global ocean covers 70 percent of our planet, makes Earth habitable, and contributes to economies, food supplies, and our health. Yet the ocean is increasingly threatened by the growing amount of carbon dioxide in the atmosphere.

Two Lamont-Doherty Earth Observatory scientists affiliated with the Center for Climate and Life are leading research projects that examine a few of the ways climate change affects the health of the ocean. Both researchers use the fossil remains of sea creatures as natural recorders of past climate and marine ecosystem changes. The information they obtain from these provides clues about how the future ocean and its inhabitants might be shaped by climate change.

Their studies are funded in part by the Center's partnership with the World Surf League PURE, which enables Lamont-Doherty scientists to pursue critical research that advances understanding of climate impacts on the ocean.

## **Ocean acidification: The other carbon dioxide problem**

Bärbel Hönisch, a marine geochemist, studies how seawater chemistry changed through time. Today, the ocean is becoming more acidic due to the rising concentration of carbon dioxide in Earth's atmosphere, about 30 percent of which is absorbed by the ocean. While this process helps to minimize global warming, the dissolution of carbon dioxide in the ocean leads to the formation of carbonic acid. As the name implies, the addition of carbonic acid makes seawater more acidic and this '[ocean acidification](#)' makes it harder for calcifying organisms such as corals, mollusks, and some plankton to build their shells and skeletons.

The current pH of the ocean is around 8.1, representing a 25 percent

increase in acidity over the past 200 years. As the amount of carbon dioxide in the atmosphere continues to rise, scientists expect seawater acidity to increase another 25 percent by the end of the 21st century. This level of acidification is similar to that of the Paleocene-Eocene Thermal Maximum (PETM), which occurred around 56 million years ago. During the PETM, a sudden rise in atmospheric carbon dioxide coincided with rapid warming and seawater acidification—conditions that lasted for 70,000 years or more.

Hönisch is analyzing the shells of tiny plankton organisms called foraminifera that were preserved in deep-sea sediments during this and other time periods, which are recovered from the sea floor by deep-ocean drilling. Her goal is to quantify changes in the ocean that occurred due to past climate shifts, and to determine whether and how [marine organisms](#) adapted to these changing conditions: Did some organisms evolve and thrive? Did some go extinct?

Hönisch is using her grant from the Center for Climate and Life to answer these questions. As part of her project, she and her research team are creating a timeline that details past levels of ocean acidity and the sensitivity of foraminifera to environmental changes. This will help them determine how past ocean temperatures and acidity affected the ability of calciferous marine organisms to build and maintain their shells. Their findings may also improve predictions of the consequences of future ecosystem changes.

"Our research has shown that the two degrees Celsius warming at the end of the last ice age had a stronger effect on foraminifer species abundance and latitudinal migration than the 0.15 unit acidification of the surface ocean," Hönisch said. "So in some ways, one might say that warming is a greater environmental stressor than acidification, at least for planktic foraminifera with this specific extent of warming and acidification. However, warming and acidification will also go hand in

hand in the future and their respective effects will add up."

Hönisch also explained that the effects of rising temperatures and acidification of seawater won't be the same throughout the [global ocean](#). "There will be a lot of variability in the conditions in the ocean," she said. "There may be refuges where certain organisms can survive."

Although observations of past ocean changes may seem grim, Hönisch noted that many marine organisms are extremely resilient to environmental change. Even the devastating impact of the [Chicxulub asteroid 66 million years ago](#) didn't completely wipe out life in the sea, so there's hope for the survival of marine organisms despite our rapidly acidifying oceans.

Hönisch's research demonstrates how information about past phenomena can be helpful today, and in the future, if we study and heed their messages, which, in this case, have been left for us in sediments at the bottom of the sea.





Scientists collect coral reef core samples in the Gulf of Chiriquí, on the Pacific coast of Panama. Credit: Brad Linsley

## **Corals: A window into past climate**

Brad Linsley, a paleoclimate scientist at Lamont-Doherty Earth Observatory, reconstructs past climate using corals and sediments to learn how changes in global temperatures, ocean salinity, and atmospheric hydrology varied in the past. He does this by analyzing microfossils preserved in deep-sea sediments and cores taken from massive corals.

The huge reef corals Linsley samples grow a skeleton at a rate of about a centimeter per year. As the skeleton grows upwards, the coral generates alternating low density and high-density band couplets at a rate of one couplet each year. These density bands are visible in X-ray images of cut slabs of the coral cores and are used by scientists to guide near-monthly resolution sampling and to help generate detailed age models.

Geochemical tracers measured in a coral's skeleton are sensitive to water temperature, salinity, river discharge, and other environmental parameters. Since healthy corals grow continuously throughout the year and can live for several centuries, massive corals can be used to create continuous records of past changes in water temperature, salinity, and other conditions that extend back several centuries.

The longest record Linsley has worked on dates back to 1521 and is from American Samoa. Other records from Panamá, Fiji, the Cook Islands, and Tonga extend back to the early 1600s. The ability to generate these detailed and accurate chronologies of past environmental conditions is what makes corals so valuable as paleoclimate archives.

Some of Linsley's recent research focuses on coral bleaching events, which are occurring with increasing frequency as the temperature of the ocean rises. Tiny algae live in the tissues of corals—they're what give corals their bright colors—and have a symbiotic relationship with the coral animals. For example, the coral provides the algae a rich supply of [carbon dioxide](#) and the algae provide corals their main source of food.



Coral cores collected by Brad Linsley and his research team in Panama. The cores will be split in half and analyzed to reconstruct the history of coral bleaching and hydrologic changes in the region back to the mid-1800s. Credit: Brad Linsley

Corals are extremely sensitive to temperature changes and when the ocean temperature rises, even one degree Celsius, corals get stressed out. When this happens, the algae are expelled from the corals, resulting in "bleached" white coral structures. During a bleaching event, algae may not disappear uniformly from coral, due in part to the fact that there may be many different species of algae present on one coral formation. In some cases, corals can recover but if the water remains warm, the corals

will usually die.

"Once a certain, site-specific temperature threshold is maintained for a specific number of weeks, many corals will bleach. However, other stresses can also cause coral death which is sometimes hard to distinguish from bleaching," Linsley said.

Coral bleaching events have occurred around the globe at the same time as some El Niño events, which cause a region of warm water to develop along the equator in the central and eastern Pacific. There may be a correlation between the two, but the exact pattern is still unclear; Linsley said that on the Pacific Ocean side of Panamá, one of his study sites, coral bleaching appears to coincide with certain very strong El Niño events.

Over the course of Linsley's career, he has studied coral bleaching events in Panamá, Fiji, and Tonga. Most recently, in March 2018, he returned to Panamá to examine a bleaching event for the second time—a trip that was supported by the Center for Climate and Life.

In Panamá, Linsley and his team collected five coral cores from an area in the Gulf of Chiriquí, on the Pacific coast. The water temperature in the region is generally 32 degrees Centigrade, or 89.6 degrees Fahrenheit, with only two degrees of seasonal variation. There's very little human development in the area, so many mangrove forests remain along the undeveloped coastline. The winds in the region move east to west and a wind shadow that blocks the tradewinds; there is both direct rainfall into the ocean and runoff from the land.

The corals record all of this environmental activity and, through different analyses, Linsley is using the cores he collected in Panamá to reconstruct the history of [coral](#) bleaching and hydrologic changes in the region back to the mid-1800s. Linsley's results will ultimately improve



understanding of seasonal and decadal-scale changes in rainfall in Central America.

This information will help the region respond to climate pressures in several ways. Agricultural business, governments, and farmers can use it to plan for future fluctuations in rainfall and implement effective crop management practices. The knowledge will also help the efforts by the Panamá Canal Authority, which operates the Panamá Canal Locks, build resilience and adapt to changes in rainfall. A large amount of water is required to move ships through the locks without the use of pumps, so a lack of rainfall could result in costly disruptions to shipping traffic through the Panamá Canal.

*This story is republished courtesy of Earth Institute, Columbia University*  
<http://blogs.ei.columbia.edu>.

Provided by Earth Institute, Columbia University

Citation: Getting warmer—understanding threats to ocean health (2018, August 16) retrieved 2 May 2024 from <https://phys.org/news/2018-08-warmerunderstanding-threats-ocean-health.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--