

Amazon mangrove forest stores twice as much carbon per acre as region's famous rainforest

September 25 2018



Credit: Oregon State University

Scientists have determined for the first time that Amazon's waterlogged coastal mangrove forests, which are being clear cut for cattle pastures and shrimp ponds, store significantly more carbon per acre than the region's famous rainforest.

The long-term study, recently published in the journal *Biology Letters*, provides a better understanding of how [mangrove](#) deforestation contributes to the greenhouse gas effect, one of the leading causes of global warming, said J. Boone Kauffman, an ecologist at Oregon State University who led the research.

The Brazilian mangrove forest fringes the entirety of the Atlantic Coast at the mouth of the Amazon, the largest river in the world with the largest mangrove forest. Although preservation of the Amazon rainforest has been the subject of intense awareness efforts over the last few decades, less attention has been paid to the Amazon mangroves.

Mangroves represent 0.6 percent of all the world's tropical forests but their deforestation accounts for as much as 12 percent of [greenhouse gas](#) emissions that come from all tropical deforestation.

"Over 25 years, we found two to three more times more carbon stored in the mangroves than in the rainforest," said Kauffman, a senior research professor in OSU's College of Agricultural Sciences. "When those forests are cut down they lose carbon, creating far more greenhouse gases than when the rainforests are cleared. Mangroves deserve conservation and participation in climate change mitigation actions throughout the world."

Mangroves are a group of trees and shrubs that live in tropical coastal intertidal zones. There are about 80 different species of mangrove trees. All of these trees grow in areas of waterlogged soils, where slow-moving waters allow fine sediments to accumulate. In these environments, mangroves sequester significant quantities of carbon that is stored for centuries.

For the study, the researchers visited nine mangroves and three salt marshes within the Brazilian Amazon. The Oregon coast, like much of

the United States, features [salt marshes](#) along its coast and estuaries. Salt marshes, mangroves and sea grass communities are collectively referred to as "blue carbon" because they hold vast [carbon](#) reservoirs and their conservation is of value with respect to climate change mitigation.

Mangrove forests are also important for biodiversity. They are known as "kindergarten of the seas." This is where young fish spawn and reproduce and spend the early parts of their lives.

Clear-cutting mangroves presents another threat: The people who live along the coast just behind these forests become more vulnerable to storms when they are cut down, Kauffman said.

"Mangroves are important for storm surge protection," he said. "When we destroy the mangroves, we make populations much more vulnerable to damage and death during hurricanes. More often these are some of the poorest people on the planet."

More information: J. Boone Kauffman et al. Carbon stocks of mangroves and salt marshes of the Amazon region, Brazil, *Biology Letters* (2018). [DOI: 10.1098/rsbl.2018.0208](https://doi.org/10.1098/rsbl.2018.0208)

Provided by Oregon State University

Citation: Amazon mangrove forest stores twice as much carbon per acre as region's famous rainforest (2018, September 25) retrieved 22 June 2024 from <https://phys.org/news/2018-09-amazon-mangrove-forest-carbon-acre.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|