

Evaluating the contribution of black carbon to climate change

September 11 2018




Fig. 1: Importance of particle size and coating amount of black carbon (BC) particles. Panel a: Particle size and number concentration can be different even if the total amount (mass) of BC is the same. Panel b: Particle size and coating amount determine the solar absorption efficiency of BC. Many BC particles are emitted as pure BC particles (BC particles without coating). BC particles are gradually coated by other aerosol species such as sulfate and organic aerosols through aerosol processes in the atmosphere (arrow). These aerosol processes enhance BC absorption efficiency by up to a factor of 2. Credit: Nagoya University

Japanese and US researchers developed an advanced model to assess the ability of black carbon particles to absorb sunlight and contribute to global warming. The model achieved higher sensitivity than obtained by previous models because it considered both particle size and the complex mixing states of black carbon in air. This advanced model will aid in the

assessment of the effectiveness of removing black carbon from the atmosphere to suppress climate change.

Black carbon refers to tiny carbon [particles](#) that form during incomplete combustion of carbon-based fuels. Black carbon particles absorb [sunlight](#), so they are considered to contribute to global warming. However, the contribution of black carbon to the heating of the Earth's [atmosphere](#) is currently uncertain. Models that can accurately assess the warming effect of black carbon on our atmosphere are needed so that we can understand the contribution of these tiny carbon particles to climate change. The mixing state of black carbon particles and their [particle size](#) strongly influence their ability to absorb sunlight, but current models have large uncertainties associated with both particle size and mixing state.

Researchers from Nagoya and Cornell Universities have combined their expertise to develop a model that can predict the direct radiative effect of black carbon with high accuracy. The team achieved such a model by considering various particle sizes and mixing states of black carbon particles in air.

"Most aerosol models are using one or two black carbon mixing states, which are not sufficient to accurately describe the mixing state diversity of black carbon in air," says Hitoshi Matsui. "Our model considers that black carbon particles have multiple mixing states in air. As a result, we can model the ability of black carbon particles to heat air more accurately than in previous estimates."




Fig 2. A schematic figure showing the treatment of particle size and coating amount in model simulations (panel a) and aerosol processes treated in the global aerosol model (panel b). Panel a: Many model simulations were made by changing particle sizes at emissions (e.g., "Small size" and "Large size"). The "Detailed method" can simulate various coating states of BC particles (pure BC, thinly-coated BC, thickly-coated BC etc.). The "Simple method", which is

similar to the method used in many previous modeling studies, cannot resolve different coating states of BC particles sufficiently. Panel b: Aerosol processes in the atmosphere (emissions, transport, transformation, and deposition processes) and their impact on BC heating effect are simulated in the global aerosol model. Credit: Nagoya University

The researchers found that the direct radiative effect of black carbon predicted by their model was highly sensitive to the particle size distribution only when the complex mixing states of black carbon were suitably described.

High sensitivity was obtained by the developed model because it calculated factors like the lifetime of black carbon in the atmosphere, the ability of black carbon to absorb sunlight, and the effect of materials coating the black carbon particles on their ability to absorb sunlight realistically. All of these factors are influenced by the particle size and mixing state of black carbon.

The results show that properly describing the particle size and mixing state of black carbon is very important to understand the contribution of black carbon to climate change.

The team's results suggest that the interactions of black [carbon](#) with atmospheric and rain patterns are likely to be more complex than previously considered. The developed model improves our ability to estimate the effectiveness of removing [black carbon](#) from the atmosphere to suppress future changes in temperature, which should help to direct research on strategies to mitigate climate change.




Fig 3. Ranges of BC heating effect (global mean). Horizontal bars in the "Detailed method" and "Simple method" show the ranges of BC heating effect when emission particle sizes are changed within their current uncertainty. The "Detailed method" has 7 times greater range of BC heating effect than the "Simple method" (0.24 W m⁻² in "Detailed method", 0.035 W m⁻² in "Simple method"). Circles in the horizontal bars show BC heating effects when average particle sizes are used for emissions. Credit: Nagoya University




Fig 4. Ratio of BC heating effect between the two simulations with the smallest and largest particle sizes at emissions. The ratio has higher values (is close to 1) when emission particle sizes are important (not important) for the estimation of BC heating effect. Credit: Nagoya University

More information: Hitoshi Matsui et al. Black carbon radiative effects highly sensitive to emitted particle size when resolving mixing-state diversity, *Nature Communications* (2018). [DOI: 10.1038/s41467-018-05635-1](https://doi.org/10.1038/s41467-018-05635-1)

Provided by Nagoya University

Citation: Evaluating the contribution of black carbon to climate change (2018, September 11) retrieved 26 April 2024 from <https://phys.org/news/2018-09-contribution-black-carbon-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.