

Efficient generation of high-density plasma enabled by high magnetic field

September 26 2018



(1) Compressing fusion fuel by nanosecond laser beams; (2) Heating the compressed fuel by a picosecond laser beam to ignite; (3) Ignition and burn Credit: Osaka University

An international joint research group led by Osaka University demonstrated that it was possible to efficiently heat plasma by focusing a relativistic electron beam (REB) accelerated by a high-intensity, shortpulse laser with the application of a magnetic field of 600 tesla (T), about 600 times greater than the magnetic energy of a neodymium magnet (the strongest permanent magnet). Their research results were published in *Nature Communications*.

If matter can be heated to temperatures of tens of millions of degrees using REB accelerated to nearly the speed of light by irradiating plasma



with high-intensity lasers, it will become possible to ignite controlled nuclear <u>fusion</u> reactions.

In the central <u>ignition</u> scheme, a prevailing scheme for inertial confinement fusion (ICF), has the problem of ignition quench, which is caused by the hot spark mixing with the surrounding cold fuel. On the other hand, in the fast ignition scheme (fast isochoric heating), a portion of low temperature fuel is heated, and then the heated region becomes the hot spark to trigger ignition before said mixing occurs. Thus, the fast ignition scheme has drawn attention as an alternative scheme.

In the fast ignition scheme, first, fusion fuel is compressed to a high density using nanosecond laser beams. Next, a high-intensity picosecond laser rapidly heats the compressed fuel, making the heated region a hot spark to trigger ignition. Nuclear fusion releases a large amount of energy by burning the majority of the fuel. (Figure 1)





Relativistic electron beam (REB) accelerated by high-intensity laser has a large divergence angle. The REB needs to be guided along the magnetic field lines to the compressed fuel core. Credit: Osaka University

The REB, which is generated by a high-intensity short-pulse laser and accelerated to nearly the speed of light, travels through high-density <u>nuclear fusion</u> fuel plasma and deposits a portion of kinetic energy in the core, making the heated region the hot spark to trigger ignition. However, REB accelerated by high-intensity lasers has a large



divergence angle (typically 100 degrees), so only a small portion of the REB collides with the core. (Figure 2)

A kilo-tesla level <u>magnetic field</u> is necessary to guide high-energy electrons at the speed of light, so the researchers employed magnetic fields of several hundreds of tesla. Because electrons, which are charged and have a small mass, easily move along a magnetic field line, they guided the high energy REB of 1MeV along the magnetic field lines to the core (the fusion fuel of 100 microns or less), achieving efficient heating of high-density plasma. They called the scheme magnetized fast isochoric heating.

In this study, laser-to-core energy coupling reached a maximum of 8 percent. The laser-to-core energy coupling, i.e., the energy deposition rate of REB, depends on the density of the plasma to be heated. In calculation based on the ignition spark formation conditions, the energy deposition rate of REB obtained in this study is several times more than that obtained by the central ignition scheme. Thus, the researchers conclude that the magnetized fast isochoric heating is very efficient and useful for the development of laser fusion energy.

Senior author Shinsuke Fujioka says, "We have made progress towards the realization of <u>laser</u> fusion energy in cooperation with researchers from home and abroad under the Joint Use/Research Center Project. Our research results will be applied to studies on the reproduction of the core of a star in laboratory simulation and the creation of new matter under extreme environments."





The energy deposition rate of REB in the core is evaluated using spectra and images of X-rays emitted from the heated plasma. Credit: Osaka University

The article, "Magnetized Fast Isochoric Laser Heating for Efficient Creation of Ultra-High-Energy-Density States," was published in *Nature Communications*.

More information: Shohei Sakata et al, Magnetized fast isochoric laser heating for efficient creation of ultra-high-energy-density states,



Nature Communications (2018). DOI: 10.1038/s41467-018-06173-6

Provided by Osaka University

Citation: Efficient generation of high-density plasma enabled by high magnetic field (2018, September 26) retrieved 3 May 2024 from <u>https://phys.org/news/2018-09-efficient-high-density-plasma-enabled-high.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.