



New kid on the block picks up relay for ozone

September 17 2018

Total ozone from Sentinel-5P



01 - 09 - 2018



The animation shows measurements of ozone for 1–13 September 2018 over Antarctica from the Copernicus Sentinel-5P satellite, showing the ozone hole over the South Pole. Sentinel-5P measurements are now being included in ESA's Climate Change Initiative to extend existing ozone records. Prior to Sentinel-5P, ozone measurements came from instruments carried on satellites such as ERS-2 and Envisat. The record goes back more than 20 years. Credit: contains modified Copernicus Sentinel data (2018), processed by DLR/BIRA

For more than 20 years, changes in ozone over Antarctica have been carefully monitored by a succession of European satellites. This important long-term record is now being added to by the Copernicus Sentinel-5P mission, which is dedicated to atmospheric monitoring.

Protecting life on Earth from the Sun's harmful rays of [ultraviolet radiation](#), the [ozone](#) layer is a very important, yet fragile, part of Earth's atmosphere.

In the 1970s and 1980s, the widespread use of damaging chlorofluorocarbons in products such as refrigerators and aerosol tins damaged ozone high up in our atmosphere. This depletion led to a hole in the ozone layer above Antarctica.


Thankfully, the 1987 Montreal Protocol on Substances that Deplete the Ozone Layer led to the phase-out and controlled use of damaging substances, which has, in turn, allowed the ozone [layer](#) to heal significantly.

The International Day for the Preservation of the Ozone Layer is marked every year on 16 September.

While the [ozone layer](#) may be healthier than it was, it is critical that it continues to be carefully monitored – not just to look back at its history, but also to see how it changes in the future.

ESA has been involved in monitoring ozone for many years. Satellites such as ERS-2 and Envisat carried instruments that measured ozone.

Total ozone Antarctica



The animation shows average ozone from 10 to 14 September for every year between 1996 and 2017 over the South Pole. ESA has been involved in monitoring ozone for many years. Satellites such as ERS-2, and Envisat carried instruments that measured ozone. These data, spanning more than two decades, are used by ESA's Climate Change Initiative to create long-term ozone records.

Credit: ESA–CCI/DLR/BIRA/KNMI

These data, along with measurements from Eumetsat's MetOp and NASA's EOS–AURA missions, span more than two decades and are used by ESA's Climate Change Initiative to create long-term records.

Now, the Copernicus Sentinel-5P satellite is adding to this vital record.

Launched in October 2017, Sentinel-5P is the first Copernicus mission dedicated to monitoring our atmosphere.

The satellite carries the state-of-the-art Tropomi instrument to map a multitude of trace gases that affect the air we breathe and our climate.

As well as measuring ozone, this new mission also measures gases such as [nitrogen dioxide](#), formaldehyde, [sulphur dioxide](#), methane, carbon monoxide, and aerosols.

Provided by European Space Agency

Citation: New kid on the block picks up relay for ozone (2018, September 17) retrieved 7 August 2024 from <https://phys.org/news/2018-09-kid-block-relay-ozone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.