

Manganese plays a key role in bacterial infection

September 20 2018



Enterococcus faecalis. Credit: Nathan Shankar, University of Oklahoma, digital colorization by Stephen Ausmus. United States Department of Agriculture, 2009.

The ability to acquire manganese during infection is essential for the virulence of *Enterococcus faecalis* in animals, according to a study published September 20 in the open-access journal *PLOS Pathogens* by José Lemos of the University of Florida College of Dentistry, and colleagues.

Manganese is an essential micronutrient for bacterial pathogens during infection. To prevent infection, the host limits manganese availability to invading bacteria through an active process known as nutritional immunity. To overcome this limitation, bacteria produce high-affinity manganese uptake systems to scavenge this nutrient from host tissues. However, little is known about the potential role of manganese in the [virulence](#) of *E. faecalis*—a leading cause of hospital-acquired infections that are often difficult to treat due to their exceptional multidrug resistance. To address this gap in knowledge, Lemos and colleagues identified and characterized the major manganese acquisition systems of this opportunistic pathogen.

The findings suggest that these transporters are essential for the growth of *E. faecalis* in manganese-restricted environments. The simultaneous inactivation of three main manganese transporters led to the loss of virulence in rabbit and mouse [infection](#) models. In contrast to the unmutated bacterial strain, the mutant strain lacking the three transporters was virtually absent in infected organs such as the heart and bladder. Collectively, the results demonstrate that manganese acquisition is essential for the virulence of *E. faecalis*. According to the authors, manganese uptake systems could be promising targets for the development of new antimicrobial therapies to combat infections.

"While we and others have shown that inactivation of certain genes and pathways can negatively affect the ability of *E. faecalis* and related bacteria to cause disease, this is the first time we have observed an outcome as dramatic as this in terms of loss of virulence in mammalian

models", notes Lemos. "In addition, this complete loss of virulence was not associated with a general loss in fitness as the mutant lacking all 3 manganese uptake systems showed normal growth and survival rates in the lab when [manganese](#) was provided in excess amounts."

More information: Cristina Colomer-Winter et al, Manganese acquisition is essential for virulence of *Enterococcus faecalis*, *PLOS Pathogens* (2018). [DOI: 10.1371/journal.ppat.1007102](https://doi.org/10.1371/journal.ppat.1007102)

Provided by Public Library of Science

Citation: Manganese plays a key role in bacterial infection (2018, September 20) retrieved 26 June 2024 from <https://phys.org/news/2018-09-manganese-key-role-bacterial-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.