


How microvilli form

September 24 2018, by Sanjay Mishra



Credit: CC0 Public Domain

The gut is lined by cells containing brush borders, which are composed of arrays of microvillar protrusions that help in nutrient absorption and provide a barrier against pathogens and toxins. Microbes such as *E. coli* can destroy microvilli with potentially life-threatening results. But how microvilli form has not been well understood.

Using super-resolution microscopy and [live cell imaging](#), Matthew Tyska, Ph.D., and colleagues show that a protein called insulin [receptor tyrosine kinase](#) substrate (IRTKS) promotes microvilli elongation. They previously discovered that IRTKS is the only protein in the intestinal brush border containing an I-BAR domain. I-BAR proteins help form membrane protrusions.

Reporting this month in the journal *Current Biology*, the researchers showed that IRTKS tracks to the distal tips of growing microvilli through the I-BAR domain. These results help explain why IRTKS is targeted by the gut pathogens that disrupt microvilli during infection of the intestinal lining.

More information: Meagan M. Postema et al. IRTKS (BAIAP2L1) Elongates Epithelial Microvilli Using EPS8-Dependent and Independent Mechanisms, *Current Biology* (2018). [DOI: 10.1016/j.cub.2018.07.022](https://doi.org/10.1016/j.cub.2018.07.022)

Provided by Vanderbilt University

Citation: How microvilli form (2018, September 24) retrieved 10 April 2024 from <https://phys.org/news/2018-09-microvilli.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--