

Burst of morning gene activity tells plants when to flower

September 24 2018



Arabidopsis thaliana plants flowering outside under natural light. Credit: Takato Imaizumi

For angiosperms—or flowering plants—one of the most important decisions facing them each year is when to flower. It is no trivial undertaking. To flower, they must cease vegetative growth and commit to making those energetically expensive reproductive structures that will

bring about the next generation.

Knowledge of this process at the cellular level is critical for understanding how plants allocate resources and produce the components we care most about—including the grains, tubers, leaves, nuts and fruits that mean so much to humans and animals alike.

In a paper published Sept. 24 in the journal *Nature Plants*, an international team of researchers has discovered that the gene FT—the primary driver of the transition to flowering in plants each spring—does something unexpected in *Arabidopsis thaliana* plants grown in natural environments, with implications for the artificial growing conditions scientists commonly used in the lab. The team, led by University of Washington biology professor Takato Imaizumi, showed that FT has a peak of [activity](#) every morning leading up to the transition, something that scientists had not previously seen in *Arabidopsis*, a model plant that is widely studied for understanding the molecular details of the transition to flowering. The morning peak of FT activity causes plants to transition earlier from [vegetative growth](#) to flowering.

"Previous research on FT activity in *Arabidopsis* showed that there is a peak of activity in the evening, not the morning," said Imaizumi, who is senior author on the paper. "We show definitively that there is a peak of morning activity—and we think we know why this morning peak was not seen previously in the research laboratory."

Prior research, which saw only an evening peak of FT gene activity, had been conducted on *Arabidopsis* plants grown indoors under fluorescent light. Imaizumi and his team—which includes researchers in Switzerland, Scotland, South Korea and Japan—grew their plants outside under sunlight.



Arabidopsis seedling. Credit: Akane Kubota

Imaizumi wanted to do this experiment because conditions at the summer solstice in Seattle, where his lab is located, are similar to the standardized, artificial "long-day" growing conditions for *Arabidopsis*: 16 hours of light and eight hours of darkness.

"I always wanted to grow plants outdoors in conditions similar to the lab just to make sure that what we're seeing in lab with *Arabidopsis* is really what's happening in nature," said Imaizumi.

His team grew non-transgenic *Arabidopsis* plants outdoors for five consecutive summers and compared them to plants grown indoors under artificial long-day conditions. Outdoor plants produced fewer leaves than indoor plants, indicating that the outdoor plants flowered earlier. Both outdoor and indoor plants showed evening peaks of FT gene activity, but outdoor-grown plants also showed a morning peak of FT activity. They concluded that the indoor, artificial growing conditions missed key qualities of natural conditions, throwing off expression of the FT gene and the trait it governs. When active, the FT gene produces a protein that travels from the leaves to the shoot apical meristem—the niche of stem cells in the shoot that produces above-ground growth—and switches the meristem from vegetative growth to floral growth.

To identify the differences between indoor and outdoor growing conditions, Imaizumi's group focused on light. The fluorescent bulbs commonly used in *Arabidopsis* research do not emit the same wavelengths of light that sunlight does. Fluorescent bulbs, for example, generate less light from far-red wavelengths. In the outdoor growing plots, the ratio of red-wavelength light to far-red wavelength was about 1-to-1, but for fluorescent bulbs this ratio is higher than 2, which means they emit more red light than far-red. When the researchers added a far-red LED lamp to the indoor growth chambers to mimic outside light, the *Arabidopsis* plants then showed a morning peak of FT gene activity.

In addition, by modifying the temperatures in the indoor growth chambers to cycle daily from about 16 degrees Celsius to almost 23 C—or from 61 degrees Fahrenheit to about 73 F—the evening FT gene activity was reduced, similar to the outdoor plants.



Arabidopsis seedlings growing outside in a Seattle summer. Credit: Akane Kubota

FT has been studied in other plants, including some crop plants, which also show morning peaks of FT expression. But most commercially important plants are too large or grow too slowly for the controlled-environment studies that are required to determine the cellular and genetic details of plant traits. That is why *Arabidopsis*, a small, fast-growing weed from the mustard family, is widely used as a substitute model organism.

The team's findings are an opportunity to revisit the artificial growing

conditions, according to Imaizumi.

"*Arabidopsis* has been studied for decades. Researchers set up their indoor growing conditions the best they could, given equipment, time and funding, and passed those conditions down to scientists they trained," said Imaizumi. "But we need to change those conditions so that what we find in the lab reflects nature more closely. If we see a change in flowering by making these minor alterations, I imagine that other traits will change as well."

Critically, their results illuminate a path forward for plant researchers to adopt artificial growth conditions that more accurately reflect natural growing conditions.

"We show that just a few simple modifications are needed to the artificial growing conditions, which researchers are using worldwide, so that lab research on *Arabidopsis* more and other plants accurately mimics outdoor growing [conditions](#)," said Imaizumi. "This ensures that the discoveries made in the lab will be more comparable to what the biological processes are—at the cellular and molecular level—in other [plants](#) of interest in nature."

More information: Young Hun Song et al, Molecular basis of flowering under natural long-day conditions in *Arabidopsis*, *Nature Plants* (2018). [DOI: 10.1038/s41477-018-0253-3](https://doi.org/10.1038/s41477-018-0253-3)

Provided by University of Washington

Citation: Burst of morning gene activity tells plants when to flower (2018, September 24) retrieved 27 April 2024 from <https://phys.org/news/2018-09-morning-gene.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.