

Probing individual edge states with unprecedented precision

September 12 2018



Measured tunneling current and its dependence on the two applied magnetic fields: The fans of red/yellow curves each correspond to a fingerprint of the conducting edge states. Each individual curve separately shows one of the edge states. Credit: University of Basel, Department of Physics

A new technique makes it possible to obtain an individual fingerprint of the current-carrying edge states occurring in novel materials such as topological insulators or 2-D materials. Physicists of the University of Basel present the new method together with American scientists in



Nature Communications.

While insulators do not conduct electrical currents, some special materials exhibit peculiar electrical properties. Though not conducting through their bulk, their surfaces and edges may support electrical currents due to quantum mechanical effects, and do so even without causing losses. Such topological insulators have attracted great interest in recent years due to their remarkable properties. In particular, their robust edge states are very promising since they could lead to great technological advances.

Similar effects as the edge states of such topological insulators also appear when a two-dimensional metal is exposed to a strong <u>magnetic</u> <u>field</u> at low temperatures. When the so-called quantum Hall effect is realized, current is thought to flow only at the edges, where several conducting channels are formed.

Probing individual edge states

Until now, it was not possible to address the numerous current-carrying states individually or to determine their positions separately. The new technique now makes it possible to obtain an exact fingerprint of the current-carrying edge states with nanometer resolution. This is reported by researchers of the Department of Physics and the Swiss Nanoscience Institute of the University of Basel in collaboration with colleagues of the University of California, Los Angeles, as well as of Harvard and Princeton University, USA.

In order to measure the fingerprint of the conducting edge states, the physicists led by Prof. Dominik Zumbühl have further developed a technique based on tunneling spectroscopy. They have used a gallium arsenide nanowire located at the sample edge which runs in parallel to the edge states under investigation. In this configuration, electrons may



jump (tunnel) back and forth between a specific edge state and the nanowire as long as the energies in both systems coincide. Using an additional magnetic field, the scientists control the momentum of tunneling electrons and can address individual edge states. From the measured tunneling currents, the position and evolution of each edge state may be obtained with nanometer precision.

Tracking the evolution

This new technique is very versatile and can also be used to study dynamically evolving systems. Upon increasing the magnetic field, the number of edge states is reduced, and their distribution is modified. For the first time, the scientists were able to watch the full edge state evolution starting from their formation at very low magnetic fields.

With increasing magnetic field, the edge states are first compressed towards the sample boundary until eventually, they move towards the inside of the sample and then disappear completely. Analytical and numerical models developed by the research team agree very well with the experimental data.

"This new technique is not only very useful to study the quantum Hall <u>edge states</u>," Dominik Zumbühl comments the results of the international collaboration. "It might also be employed to investigate new exotic materials such as <u>topological insulators</u>, graphene or other 2-D materials."

More information: T. Patlatiuk et al, Evolution of the quantum Hall bulk spectrum into chiral edge states, *Nature Communications* (2018). DOI: 10.1038/s41467-018-06025-3



Provided by University of Basel

Citation: Probing individual edge states with unprecedented precision (2018, September 12) retrieved 1 May 2024 from https://phys.org/news/2018-09-probing-individual-edge-states-unprecedented.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.