

Scientists propose a new method for detecting dangerous nitrogen-containing liquids

September 11 2018



NMR spectrum of mmonium nitrate. Credit: Galina Kupriyanova

A team of researchers from the Institute of Physical and Mathematical Sciences and Information Technologies of Immanuel Kant Baltic Federal University (BFU), together with their colleagues from Gebze Technical University, used nuclear magnetic resonance to detect toxic and flammable nitrogen-containing liquids. The study was published in *Chemical Physics*.

Nuclear magnetic resonance (NMR) is the effect of energy absorption or



emission by a substance that contains nuclei with nonzero spin in its external magnetic field. The set of resonance lines is individual for each substance and is used for identification. The stronger the magnetic field, the more information researchers can determine about substance structure. The most intensive NMR signals are observed in hydrogen nuclei.

The physicists suggested a new method for detecting explosives and toxic <u>nitrogen</u>-containing liquids based on the NMR relaxation method in a low magnetic field on nitrogen nuclei ¹⁴N. They also developed a protocol of measurements that can be used in security screening procedures. According to the scientists, traditional ¹H NMR method cannot be used at a checkpoint because the equipment is too massive and expensive, and the analysis takes too long. The method based on studying ¹⁴N NMR relaxation signals (NMR relaxometry) provides for a different technical approach reducing installation and operation costs and speeding up analytical processes. Another advantage of the suggested approach is the analysis of ¹⁴N NMR relaxation signals which do not depend on the intermolecular interaction and only slightly depend on the concentration of the substance.

In their work, the scientists measured the ¹⁴N NMR relaxation times in liquids. They analyzed three groups of nitrogen-containing substances: explosive and flammable liquids, compounds with a nitrate group (nitrogen bound with three oxygen atoms); and toxic substances with a nitrogen atom not bound to oxygen. As a result, the scientists discovered a range of relaxation time values typical for explosives and toxic substances. This range can be used for their identification.

"In this work we showed that ¹⁴N NMR is an efficient method for detecting large quantities of dangerous nitrogen-containing substances. We solved a number of technical issues associated with low resonance frequency and broad NMR spectrum bands. We also succeeded in



successfully detecting ¹⁴N NMR signals from different nitrogen-based substances and obtaining signal relaxation parameters. The majority of toxic and flammable liquids measured in our work are characterized by short <u>relaxation</u> times. This is how they can be separated from other relatively safe materials. An important advantage of the suggested NMR relaxometry method is the short time required to detect dangerous substances," says Professor Galina Kupriyanova, the author of the study, and doctor of physics and mathematics.

According to the researchers, portable modern NMR relaxometry devices are precise enough to detect dangerous substances at checkpoints.

More information: G.V. Mozzhukhin et al, Low-field ¹⁴N nuclear magnetic resonance for detection of dangerous liquids, *Chemical Physics* (2018). DOI: 10.1016/j.chemphys.2018.07.032

Provided by Immanuel Kant Baltic Federal University

Citation: Scientists propose a new method for detecting dangerous nitrogen-containing liquids (2018, September 11) retrieved 2 May 2024 from <u>https://phys.org/news/2018-09-scientists-method-dangerous-nitrogen-containing-liquids.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.