

Three new species of fish discovered in the extreme depths of the Pacific Ocean

September 10 2018



CT scan of the Atacama snailfish. Credit: Newcastle University

An exploration to one of the deepest places on earth has captured rare footage of what is believed to be three new species of the elusive Snailfish.

Involving a team of 40 scientists from 17 different nations, including Dr. Alan Jamieson and Dr. Thomas Linley from Newcastle University, UK, the expedition to the Atacama Trench has uncovered a wealth of



information about life in one of the deepest places on earth.

The pink, the blue and the purple

Among the new discoveries are what the team believe to be three new species of snailfish.

Temporarily named 'the pink, the blue and the purple Atacama Snailfish', the footage shows the fish feeding and interacting in their secret world 7,500 metres below the surface.

These fish are part of the Liparidae family and do not conform to the preconceived stereotypical image of what a deep-sea fish should look like.

Instead of giant teeth and a menacing frame, the fishes that roam in the deepest parts of the ocean are small, translucent, bereft of scales—and highly adept at living where few other organisms can. The snailfish will be featured as part of the Challenger Conference 2018 which kicks off at Newcastle University from today and runs until Friday.

Perfectly adapted to living at extreme pressure

Dr. Thomas Linley, from Newcastle University, said:

"There is something about the snailfish (fish of the family Liparidae) that allows them to adapt to living very deep. Beyond the reach of other fish they are free of competitors and predators.

"As the footage clearly shows, there are lots of invertebrate prey down there and the snailfish are the top predator, they seem to be quite active and look very well-fed.



"Their gelatinous structure means they are perfectly adapted to living at <u>extreme pressure</u> and in fact the hardest structures in their bodies are the bones in their inner ear which give them balance and their teeth. Without the extreme pressure and cold to support their bodies they are extremely fragile and melt rapidly when brought to the surface."

Amazingly, the team did manage to catch one the <u>new species</u> of snailfish which followed its amphipod prey into one of the traps. The single specimen was in very good condition and, following careful preservation, is currently being described by the Newcastle team with the help of colleagues from the United States and the Natural History Museum, London.

Pioneering technology for exploration of the ultradeep

The Hadal Trenches are one of the last great frontiers in marine science and the deepest places on Earth.

Mostly located around the Pacific rim in areas where tectonic plate collide and plunge, the seafloor reaches depths close to 11,000 metres (~7miles) in some areas.





The Atacama snailfish. Credit: Newcastle University

The Atacama Trench, a trench almost 6000 km long and more than 8000 m deep, runs along the west coast of South America.

Newcastle University scientists and engineers have been pioneering technology for the exploration of these ultra-deep environments for the last five years and have to date completed nearly 250 deployments of their novel 'lander' systems.

Using two full-ocean depth (11,000 m) capable landers equipped with HD cameras and traps the Newcastle team assessed the animals found within the trench.

Designed and developed at Newcastle University, the lander is dropped



overboard, and free-falls to the ocean floor where it carries out a variety of monitoring and sampling tasks.

It can take four hours for a trap to sink to the bottom and after waiting an additional 12 to 24 hours, the researchers send an acoustic signal to the trap, which releases weights and the lander rises to the surface with the help of floatation. This allows the team to catch fish specimens and take video footage of life at the bottom of the ocean.

On this latest expedition to the Atacama Trench in the SE Pacific off the coast of Peru and Chile, the team deployed their baited camera system 27 times from 2537 to the deepest point, Richard's Deep, at just over 8000 metres. More than 100 hours of video and 11,468 photographs were taken at the seabed.

Somersaulting isopod

As well as the snailfish, the team also filmed some astonishingly rare footage of long-legged isopods, known as Munnopsids, which are about the size of an adult hand. These crustaceans have small bodies, extraordinarily long legs and swim backwards and upsides down, propelling themselves with paddles on their ventral side – their 'tummies' - before righting themselves on the seafloor and spreading their long walking legs out like a spider.

"We don't know what species of munnopsid these are but it's incredible to have caught them in action in their natural habitat—especially the flip they do as they switch from swimming to walking mode," says Thom.

Provided by Newcastle University



Citation: Three new species of fish discovered in the extreme depths of the Pacific Ocean (2018, September 10) retrieved 27 June 2024 from <u>https://phys.org/news/2018-09-species-fish-extreme-depths-pacific.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.