

Tracking hydrogen movement using subatomic particles

September 26 2018



Figure: When a negative muon $(\mu$ -) is implanted into MgH2, the μ - is trapped by the muon atomic orbitals near a Mg nucleus. Since μ - has a polarized spin, we can obtain information about the magnetic field at the Mg nucleus sites, formed by the hydrogen nucleuses, through the observation of how the μ - spin depolarizes with time. Credit: J. Sugiyama/Toyota Central Research and Development Laboratories Inc.

A muon is an unstable subatomic particle similar to an electron but with much greater mass. The lifetime of a muon is only a couple of microseconds, but this is long compared with the lifetimes of many



unstable subatomic particles. Because of their comparatively long lifetime, positive muons are often used to detect internal magnetic fields in solid materials. However, negative muons have seldom been used for this purpose because a large data set is required to obtain reliable results and experimental data collection times are normally limited. Recently, researchers developed a system that can count muon events at a much faster rate, allowing an experiment to be completed in a suitable time frame. Using this system, a Japanese collaboration has realized the longstanding goal of using negative muons to observe the local nuclear magnetic fields in a solid for the first time.

As explained in their article in *Physical Review Letters*, the team used magnesium hydride as the solid in their experiments. Magnesium hydride has a formula of MgH_2 and is a potential candidate as a hydrogen storage material. Magnesium hydride was selected for study in experiments using the negative muon beam because muons initially captured on hydrogen are transferred quickly to magnesium, which allowed the transfer process of hydrogen to be investigated.

"The magnesium atoms exposed to the negative muon beam were effectively converted to sodium," says first author Jun Sugiyama at Toyota Central Research & Development Laboratories Inc. "The local <u>magnetic field</u> of the hydrogen atoms around these converted atoms was then able to be detected, which meant that we could track hydrogen diffusion."

The experiments used a high-intensity muon beam and highly integrated positron detector system to detect the local magnetic fields in the magnesium hydride sample. The obtained spectra were consistent with the magnesium atoms having a random magnetic field, agreeing with theoretical predictions. In particular, the results agreed with estimations from dipole field calculations, indicating that the nuclear magnetic fields of hydrogens in <u>magnesium</u> hydride were indeed observed.



"Our approach using negative muons to detect the local behavior of ions is attractive because it allows us to study the dynamics of light elements in a solid from the fixed point of the nucleus," says co-author Kazuhiko Ninomiya at Osaka University. "This approach is therefore complementary to nuclear magnetic resonance spectroscopy."

Using this negative muon-based technique, it is now possible to track the movement of <u>hydrogen</u> in a solid, which should aid the development of <u>hydrogen storage materials</u>.

More information: Jun Sugiyama et al. Nuclear Magnetic Field in Solids Detected with Negative-Muon Spin Rotation and Relaxation, *Physical Review Letters* (2018). DOI: 10.1103/PhysRevLett.121.087202

Provided by Osaka University

Citation: Tracking hydrogen movement using subatomic particles (2018, September 26) retrieved 10 May 2024 from <u>https://phys.org/news/2018-09-tracking-hydrogen-movement-subatomic-particles.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.