

Virus may combat fire ants, but caution is needed

September 12 2018



A group of red imported fire ant workers feed on a meal worm. Credit: Kyoto University/Yang

Native to South America, imported red fire ants were introduced accidentally into the United States in the early 20th century. These ants subsequently invaded other countries, including Australia, New Zealand,

Taiwan, China, and more recently, Japan and South Korea.

A survey in Japan and Korea has revealed that fire ants are mostly confined to areas in or near container yards or warehouses. However, a Kyoto University study published in *Scientific Reports* suggests that results of this survey may need to be interpreted cautiously, as conventional [food](#) lures may result in underestimating actual ant numbers.

Fire ants are known to love oily foods, and hence potato chips or hot dog slices are often used in surveys in the United States, Australia, and Taiwan. The Kyoto team discovered that ants infected with a particular [virus](#) forage much less and prefer carbohydrates such as diluted honey over oily and protein-rich foods like tuna and peanut butter.

"This finding might lead to the impression that the pathogen can help us combat [invasive ants](#), but it hinges on our understanding of the role of that virus in the ant's biology," says corresponding author Chin-Cheng Scotty Yang of Kyoto University.

Yang elaborates that the lack of fire ants at a lure station doesn't mean that fire ants are absent. It could be a result of reduced foraging or alterations in food preference induced by the virus.

"These findings also suggest that viral prevalence in the field may potentially influence the efficacy of conventional chemical [control](#) by low-toxic baits," continues co-author DeWayne Shoemaker at the University of Tennessee.

Use of low-toxicity baits for ant control requires that the ants consume a sufficiently lethal quantity. Shoemaker explains that this particular virus appears to disrupt fire ant foraging and food preferences, raising concerns that infected ants may not ingest a sufficient dose of bait for

successful control.

"We are now considering how to integrate the virus into existing chemical control schemes," continues Yang. The team anticipates that this could lead to more efficient and effective [fire ant](#) control.

The study includes contributions from a multi-institutional invasive ant management intervention consortium supported by the Kyoto University Research Coordination Alliance program, and includes research groups from Japan, Taiwan, and the United States.

One goal of this consortium is to better understand the biology and interactions of invasive ants and their pathogens, to mitigate their impact.

More information: Hung-Wei Hsu et al, Viral infections in fire ants lead to reduced foraging activity and dietary changes, *Scientific Reports* (2018). [DOI: 10.1038/s41598-018-31969-3](https://doi.org/10.1038/s41598-018-31969-3)

Provided by Kyoto University

Citation: Virus may combat fire ants, but caution is needed (2018, September 12) retrieved 15 June 2024 from <https://phys.org/news/2018-09-virus-combat-ants-caution.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|