

Wetlands are key for accurate greenhouse gas measurements in the Arctic

September 11 2018



Western Russian tundra is a mosaic of dry and wet ecosystem types, functioning very differently with respect to carbon exchange. Credit: Maija Marushchak

The Arctic is rapidly warming, with stronger effects than observed elsewhere in the world. The Arctic regions are particularly important with respect to climate change, as permafrost soils store huge amounts of the Earth's soil carbon (C). Warming of Arctic soils and thawing of permafrost can have substantial consequences for the global climate, as the large C stored in soils could be released to the atmosphere as the greenhouse gases carbon dioxide (CO₂) and methane (CH₄). The release of these heat-trapping gases, in turn, has the potential to further enhance climate warming.

Determining whether the Arctic is continuing to take up carbon from the atmosphere or instead releasing it to the atmosphere is an urgent research priority, particularly as the climate warms. A new study by researchers at the University of Eastern Finland now provides the first estimate of regional carbon budget for tundra in Western Russia for the 10-year period from 2006 to 2015. The researchers found that over the past decade, the region has likely remained a net carbon sink, sequestering atmospheric CO₂ through plant uptake and growth. This signal varied little between all the years and was particularly strong in wetlands, which were "hotspots" for carbon uptake. Wetlands are also "hotspots" of methane emissions in the region, making the identification of wetlands essential for determining the regional carbon budget. However, it remains challenging to determine the area of tundra wetlands at broader scales because they can be difficult to identify from satellite images, requiring many measurements on the ground to verify their locations.

Due to harsh winter conditions, making measurements throughout the year in tundra sites is exceptionally difficult. Few measurements have been made, making the assessment of the Arctic carbon balance challenging. Previously, researchers from the Biogeochemistry research group at the University of Eastern Finland have measured carbon balance in Western Russia during four growing seasons, but also during

more rarely studied periods the spring, fall and winter, laying the groundwork for a longer-term assessment of whether the site has continued to take up atmospheric carbon. Now, the researchers from this group are evaluating just that: is this region a net [carbon](#) sink over a decade and why or why not?

"In many major research projects, a synthesis of the data collected by using models is always left to be done in the last phase. Owing to time limitations, this important task generally remains unaccomplished. To that end, we are proud that this work signifies an important achievement," Dr Narasinha Shurpali from the University of Eastern Finland says.

The results, based on model simulations with various ecosystem models, were recently published in *Global Change Biology* – a leading journal in environmental science.

More information: Claire C. Treat et al. Tundra landscape heterogeneity, not interannual variability, controls the decadal regional carbon balance in the Western Russian Arctic, *Global Change Biology* (2018). [DOI: 10.1111/gcb.14421](https://doi.org/10.1111/gcb.14421)

Provided by University of Eastern Finland

Citation: Wetlands are key for accurate greenhouse gas measurements in the Arctic (2018, September 11) retrieved 3 May 2024 from <https://phys.org/news/2018-09-wetlands-key-accurate-greenhouse-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
