



nuclei during the Big Bang are crucial in the determining the primordial abundance of lithium, the third (and last) chemical element formed during the very early phase of the creation of the universe. The standard models of the Big Bang predict an abundance of  $\text{Li}^7$ , the main lithium isotope, three or four times more than that actually observed. Recently, at the n\_TOF facility at CERN, researchers investigated the possibility of a [neutron](#) channel that might be able to increase the destruction rate of the isotope  $\text{Be}^7$ , the precursor of  $\text{Li}^7$ , and therefore make the calculated and observed cosmological abundance of lithium compatible.

"Potentially, a neutron reaction channel might be able to resolve the cosmological lithium problem, which is one of the still unresolved aspects of the current standard description of the Big Bang," says University of Seville Professor José Manuel Quesada.

At the SINQ facility at PSI (Villigen Switzerland) the "uncut" material for use in the new experiment was separated. The material was then sent to the ISOLDE radioactive beam facility at CERN to produce a pure target with less than 0.1 milligrams of  $\text{Be}^7$ , which was then sent to the n\_TOF facility to be included in neutron measurements.

This is the first time that the two CERN facilities dedicated to nuclear physics experiments have carried out an experiment together, using the ISOLDE radioactive ion beam to produce the target necessary for an experiment at n\_TOF using the neutron time-of-flight technique.

In a previous experiment at n\_TOF, the effective section of the  $^7\text{Be}(n,\alpha)^4\text{He}$  reaction was measured in a wide range of energies, which allowed for the imposition of strict restrictions on one of the destruction mechanisms of isotope  $\text{Be}_7$  during the Big Bang. In this experiment, however, the reaction  $^7\text{Be}(n,p)^7\text{Li}$  was measured, extending previously acquired data to a greater range of energies, allowing for the updating of the [reaction](#) rate used in the calculations in the standard model of the Big

Bang.

"Although the new data obtained from the experiments at n\_TOF allow for the establishment of a much firmer basis for BBN calculations, the conclusion of this project is that neutron channels are not enough to resolve the cosmological [lithium](#) problem. The scientific community has a challenge that will require additional efforts to resolve, and this will involve the fields of nuclear astrophysics, astronomic observations, non-standard cosmology and even new physics beyond the Standard Model of particle physics," the researchers write.

**More information:** L. Damone et al,  $\text{Be}^7(\text{n,p})\text{Li}^7$  Reaction and the Cosmological Lithium Problem: Measurement of the Cross Section in a Wide Energy Range at n\_TOF at CERN, *Physical Review Letters* (2018). [DOI: 10.1103/PhysRevLett.121.042701](https://doi.org/10.1103/PhysRevLett.121.042701)

Provided by University of Seville

Citation: The cosmological lithium problem (2018, October 10) retrieved 2 May 2024 from <https://phys.org/news/2018-10-cosmological-lithium-problem.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---