


Harvesting solar fuels through a bacterium's unusual appetite for gold

October 9 2018, by Theresa Duque



A single nanocluster of 22 gold atoms -- Au₂₂ -- is only 1 nanometer in diameter, allowing it to easily slip through the bacterial cell wall. Credit: Peidong Yang, UC Berkeley

A bacterium named *Moorella thermoacetica* won't work for free. But UC Berkeley researchers have figured out it has an appetite for gold. And in exchange for this special treat, the bacterium has revealed a more efficient path to producing solar fuels through artificial photosynthesis.

M. thermoacetica first made its debut as the first non-photosensitive bacterium to carry out [artificial photosynthesis](#) in a study led by Peidong Yang, a professor in UC Berkeley's College of Chemistry. By attaching light-absorbing nanoparticles made of cadmium sulfide (CdS) to the bacterial membrane exterior, the researchers turned M. thermoacetica into a tiny photosynthesis machine, converting sunlight and carbon dioxide into useful chemicals.

Now Yang and his team of researchers have found a better way to entice this CO₂-hungry bacterium into being even more productive. By placing light-absorbing gold nanoclusters inside the bacterium, they have created a biohybrid system that produces a higher yield of chemical products than previously demonstrated. The research, funded by the National Institutes of Health, was published on Oct. 1 in *Nature Nanotechnology*.

For the first hybrid model, M. thermoacetica-CdS, the researchers chose cadmium sulfide as the semiconductor for its ability to absorb [visible light](#). But because cadmium sulfide is toxic to bacteria, the nanoparticles had to be attached to the cell membrane "extracellularly," or outside the M. thermoacetica-CdS system. Sunlight excites each [cadmium-sulfide](#) nanoparticle into generating a charged particle known as an electron. As these light-generated electrons travel through the bacterium, they interact with multiple enzymes in a process known as "CO₂ reduction," triggering a cascade of reactions that eventually turns CO₂ into acetate, a valuable chemical for making [solar fuels](#).

But within the extracellular model, the electrons end up interacting with other chemicals that have no part in turning CO₂ into acetate. And as a result, some electrons are lost and never reach the enzymes. So to improve what's known as "quantum efficiency," or the [bacterium's](#) ability to produce acetate each time it gains an electron, the researchers found another semiconductor: nanoclusters made of 22 gold atoms (Au₂₂), a material that M. thermoacetica took a surprising shine to.

"We selected Au₂₂ because it's ideal for absorbing visible light and has the potential for driving the CO₂ reduction process, but we weren't sure whether it would be compatible with the bacteria," Yang said. "When we inspected them under the microscope, we discovered that the bacteria were loaded with these Au₂₂ clusters—and were still happily alive."

Imaging of the *M. thermoacetica*-Au₂₂ system was done at UC Berkeley's Molecular Imaging Center.

The researchers also selected Au₂₂ — dubbed by the researchers as "magic" gold nanoclusters—for its ultras-small size: A single Au₂₂ nanocluster is only 1 nanometer in diameter, allowing each nanocluster to easily slip through the bacterial cell wall.

"By feeding bacteria with Au₂₂ nanoclusters, we've effectively streamlined the electron transfer process for the CO₂ reduction pathway inside the bacteria, as evidenced by a 2.86 percent quantum efficiency—or 33 percent more acetate produced within the *M. thermoacetica*-Au₂₂ system than the CdS model," Yang said.

The magic gold nanocluster is the latest discovery coming out of Yang's lab, which for the past six years has focused on using biohybrid nanostructures to convert CO₂ into useful chemicals as part of an ongoing effort to find affordable, abundant resources for renewable fuels, and potential solutions to thwart the effects of climate change.

"Next, we'd like to find a way to reduce costs, improve the lifetimes for these biohybrid systems, and improve quantum efficiency," Yang said. "By continuing to look at the fundamental aspect of how gold nanoclusters are being photoactivated, and by following the electron transfer process within the CO₂ reduction pathway, we hope to find even better solutions."

More information: Hao Zhang et al, Bacteria photosensitized by intracellular gold nanoclusters for solar fuel production, *Nature Nanotechnology* (2018). [DOI: 10.1038/s41565-018-0267-z](https://doi.org/10.1038/s41565-018-0267-z)

Provided by University of California - Berkeley

Citation: Harvesting solar fuels through a bacterium's unusual appetite for gold (2018, October 9)
retrieved 10 April 2024 from
<https://phys.org/news/2018-10-harvesting-solar-fuels-bacterium-unusual.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
