

ICESat-2 laser fires for first time, measures Antarctic height

October 4 2018, by Kate Ramsayer




Illustration of ICESat-2. Credit: NASA's Goddard Space Flight Center

The laser instrument that launched into orbit last month aboard NASA's Ice, Cloud and land Elevation Satellite-2 (ICESat-2) fired for the first time Sept. 30. With each of its 10,000 pulses per second, the instrument is sending 300 trillion green photons of light to the ground and measuring the travel time of the few that return: the method behind ICESat-2's mission to monitor Earth's changing ice. By the morning of


Oct. 3, the satellite returned its first height measurements across the Antarctic ice sheet.

"We were all waiting with bated breath for the lasers to turn on and to see those first photons return," said Donya Douglas-Bradshaw, the project manager for ICESat-2's sole instrument, called the Advanced Topographic Laser Altimeter System, or ATLAS. "Seeing everything work together in concert is incredibly exciting. There are a lot of moving parts and this is the demonstration that it's all working together."

ICESat-2 launched on Sept. 15 to precisely measure heights and how they change over time. It does this by timing how long it takes individual photons to leave the satellite, reflect off the surface, and return to receiver telescope on the satellite. The ATLAS instrument can time photons with a precision of less than a billionth of a second, which allows the mission to detect small changes in the planet's ice sheets, glaciers and sea ice.

Once ICESat-2 was in space, the ATLAS team waited to turn on the lasers for about two weeks to allow any Earthly contaminants or gases to dissipate.

"It's very critical when you fire the lasers that you don't have contaminants because you could damage the optics," Douglas-Bradshaw said. "Fourteen days is well beyond the time needed for that, but we wanted to be safe."



A visualization of ICESat-2 data, called a photon cloud, shows the first set of height measurements from the satellite, taken as it orbited over the Antarctic ice sheet. Each blue dot represents a photon detected by the ATLAS instrument. This photon cloud shows the elevation measured by photons in the middle of the ice sheet, following along 6.2 miles (10 kilometers) of the satellite's ground track, from left to right. The speckled dots are background photons from sunlight, but the thick blue line is actually a concentration of dots that represent laser photons that returned to the ICESat-2 satellite. Credit: NASA's Goddard Space Flight Center

During those two weeks, the ICESat-2 operations team turned on and tested the various systems and subsystems of the spacecraft and instrument, and fired thrusters to start placing the satellite in its final polar orbit, approximately 310 miles (500 kilometers) above Earth.

Before the [laser](#) was even turned on, however, the team eagerly awaited another milestone, Douglas-Bradshaw said. The door that protected the telescope and detector elements during launch had to be opened. The team had two chances to release one of two spring-loaded pins to open the door. This was successfully accomplished on Sept. 29.

The following day, it was the laser's turn. The engineering team had been working with the operations team that controls the instrument on orbit, so the commands were ready to go—first turning on the laser itself, waiting for it to warm up, and then issuing commands to put it in fire mode.

The laser energy levels jumped up, and the device that starts ATLAS's sophisticated stopwatch was active—two different, independent indicators that the laser was firing away.

"We were all incredibly excited and happy, everyone was taking pictures of the screens showing data plots," Douglas-Bradshaw said. "Someone noted: 'Now we have a mission, now we have an instrument.'"

Three days later, the ICESat-2 team had the first segment of height data, taken as the satellite flew over Antarctica.

Computer programmers were up all night analyzing the latitude, longitude and elevation represented by each [photon](#) that returned to the ATLAS instrument—and by 6 a.m., Tom Neumann, ICESat-2 deputy project scientist, was texting screenshots of the height data to the rest of the team.

"It was awesome," Neumann said. "Having it in space, and not just simulating data on the ground, is amazing. This is real light that went from ATLAS to Earth and back again."

When scientists analyze the preliminary ICESat-2 data, they examine what is called a "photon cloud," or a plot of each photon that ATLAS detects. Many of the points on a photon cloud are from background photons—natural sunlight reflected off Earth in the exact same wavelength as the laser photons. But with the help of computer programs that analyze the data, scientists can extract the signal from the noise and

identify height of the ground below.

The first photon cloud generated by ICESat-2 shows a stretch of elevation measurements from East Antarctica, passing close to the South Pole at a latitude of 88 degrees south, then continuing between Thwaites Glacier and Pine Island Glacier in West Antarctica.

Next up for ICESat-2 is a suite of procedures to optimize the [instrument](#), Neumann said, including tests to ensure the laser is pointing at the precisely correct angle and lasing at the precisely correct wavelength to allow as many photons as possible to hit the detector.

"It will take a couple of additional weeks," he said, "but about one month after launch we'll hopefully start getting back some excellent science-quality data."

Provided by NASA's Goddard Space Flight Center

Citation: ICESat-2 laser fires for first time, measures Antarctic height (2018, October 4)
retrieved 20 March 2024 from <https://phys.org/news/2018-10-icesat-laser-antarctic-height.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
