

## Marine algae bloom-derived biotoxins alter the development of zooplankton and the ocean food web

October 24 2018



Credit: CC0 Public Domain

Researchers from the University of Barcelona in collaboration with a



team from the Zoological Station Anton Dohrn (Italy) have analysed the impact of diatom algae in the development of Oikopleura dioica (O. Dioica), a type of marine zooplankton invertebrate that plays an important role in the global dynamics of the marine food webs and the biosphere carbon cycle. The results, published in *Communications Biology*, show that biotoxins that are produced by these algae at the end of the blooms can affect embryonic development, and therefore the reproduction of O. Dioica, with severe ecological consequences.

The study is especially relevant, considering that ocean acidification and warming caused by climate change could intensify the frequency of blooms of <u>harmful algae</u> like diatoms.

Diatoms are a type of microalgae that contribute to marine phytoplankton photoautotrophic production, since they provide nutrients to all other forms of life without of solar light. However, diatoms can produce biotoxins (polyunsaturated aldehydes) as a defence mechanism against predators such as copepods. These substances are massively released at the end of the blooms and can alter the biology of marine animals in the area.

The new study has analysed the impact of these biotoxins in the development of O. dioca, a type of marine invertebrate of the appendicularian chordates (Appendicularia or Larvacea). Appendicularians are key organisms in the ocean food web, since they process about 10 percent of the primary production in the ocean and serve as food for fish larvae in the marine trophic chain.

"Results show that biotoxins that are produced by diatoms can cause important alterations in the development of O. dioica inducing a phenotype they call 'golf ball,' which affects the embryo morphogenesis and differentiation of their trunk and tail," says researcher Cristian Cañestero. "Although in our study, embryonic malformations appear in



higher concentrations than the ones found in natural conditions, we have also observed lethal embryonic anomalies at similar concentrations to the measured values in nature after blooms," says the researcher.

This discovery is especially relevant considering the increase of acidification and warming in the oceans due climate change could intensify the seriousness and frequency of algae blooms. "Discovering how blooms in harmful algae and the massive production of toxins can alter the physiology of larvae is important due its impact in the marine food web and to predict future problems in fishing caused by global climate change," notes the researcher.

## A potential biomarker of zooplankton

The study also analysed the genic response of O. dioica to the impact of biotoxins. "We found that O. dioica can respond to the stress caused by diatoms by delaying the zygotic transcription of the development genes during the first stages and activating defensive genes (set of genes that protect the cell against environmental attacks), even in innocuous concentrations of <u>biotoxins</u> in which we did not observe any embryonic alteration," says Cristian Cañestero.

According to the conclusions of the study, this response could be part of a mechanism developed through evolution that protects embryos from environmental dangers and which would serve as a potential biomarker. "Defensive genes would be a molecular biosensor that marine ecologists could use to control the genetic stress in natural populations of appendicularians, and other organisms, exposed to <u>algae blooms</u> in their habitat, and afterwards, assess the potential impact of blooms associated with climate change," says the researcher.

These results open a new study area for the UB research team. "The article brings the basics of a new research line in the EcoEvoDevo field,



the intersection between ecology, developmental biology and evolution, which will do research on whether the embryos of marine organisms are ready to respond to <u>climate change</u>," concludes Cristian Cañestero.

**More information:** Nuria P. Torres-Águila et al. Diatom bloomderived biotoxins cause aberrant development and gene expression in the appendicularian chordate Oikopleura dioica, *Communications Biology* (2018). <u>DOI: 10.1038/s42003-018-0127-2</u>

Provided by University of Barcelona

Citation: Marine algae bloom-derived biotoxins alter the development of zooplankton and the ocean food web (2018, October 24) retrieved 3 May 2024 from https://phys.org/news/2018-10-marine-algae-bloom-derived-biotoxins-zooplankton.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.