


Are we losing one of our biggest carbon dioxide sinks?

November 1 2018



Eelgrass blue carbon stocks compared to other marine and terrestrial ecosystems.

Data sources: Kennedy and Björk, 2009, Fourqurean et al., 2012, Duarte et al., 2013, Röhr et al., 2018. Credit: © Emilia Röhr/Åbo Akademi University.

In a new study spanning coastal areas of the Northern Hemisphere, a coordinated research network led by MSc Emilia Röhr, Assoc. Prof. Christoffer Boström from Åbo Akademi University and Prof. Marianne Holmer from University of Southern Denmark explored the magnitude of organic carbon stocks stored and sequestered by eelgrass (*Zostera marina*) meadows—the most abundant seagrass species in temperate waters.

"We discovered that eelgrass organic [carbon](#) stocks were comparable to organic carbon stocks of tropical seagrass species, as well as mangroves, salt marshes and terrestrial ecosystems," Emilia Röhr says.

On average, [eelgrass meadows](#) stored 27.2 tons of organic carbon per hectare, although the variation between the regions was considerable (from three to 265 tons per hectare). Hotspots for carbon sequestration were identified especially in the Kattegat-Skagerrak region and southern parts of the Baltic Sea, where the organic carbon stocks were over eight times higher than in the Archipelago Sea of Finland.

The high carbon storage capacity of [eelgrass](#) meadows argues for protection and restoration of this unique ecosystem. The areas with the highest carbon stock capacity especially deserve recognition as part of global carbon marketing programmes.

"Terrestrial forests are well known for their capacity to store carbon (green carbon), while the so-called blue carbon stored and sequestered by coastal vegetated ecosystems, such as mangroves, saltmarshes, macroalgae and seagrasses, have received much less attention," says

Röhr.



Eelgrass, *Zostera marina*. Credit: © Christoffer Boström/Åbo Akademi University

"Although these cover only 0.5 percent of the seafloor, their carbon storage capacity accounts for more than 55 percent of the carbon stored by photosynthetic activity on Earth."

In the marine systems, the blue carbon species alone accounts for up to 33 percent of the total oceanic CO₂ uptake. In contrast to terrestrial soils, which usually store carbon for decades, the carbon stored in blue carbon ecosystems may persist for timescales of millennia or longer, and thus contribute significantly to climate change mitigation and alleviation of the rising atmospheric carbon dioxide concentrations. Despite the importance of these ecosystems, to date, none of them are included in the global carbon trading programmes. Alarmingly, in the past 50 years, at least one-third of the distribution area of coastal vegetated [ecosystems](#) has been lost.



Eelgrass, *Zostera marina*. Credit: © Christoffer Boström/Åbo Akademi University

More information: Maria Emilia Röhr et al, Blue Carbon Storage Capacity of Temperate Eelgrass (*Zostera marina*) Meadows, *Global Biogeochemical Cycles* (2018). [DOI: 10.1029/2018GB005941](https://doi.org/10.1029/2018GB005941)

Provided by Åbo Akademi University

Citation: Are we losing one of our biggest carbon dioxide sinks? (2018, November 1) retrieved 6 May 2024 from <https://phys.org/news/2018-11-biggest-carbon-dioxide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.