


One step closer to complex quantum teleportation

November 5 2018



Credit: CC0 Public Domain

The experimental mastery of complex quantum systems is required for future technologies like quantum computers and quantum encryption. Scientists from the University of Vienna and the Austrian Academy of

Sciences have broken new ground. They sought to use more complex quantum systems than two-dimensionally entangled qubits and thus can increase the information capacity with the same number of particles. The developed methods and technologies could in the future enable the teleportation of complex quantum systems. The results of their work, "Experimental Greenberger-Horne-Zeilinger entanglement beyond qubits," is published recently in the renowned journal *Nature Photonics*.

Similar to bits in conventional computers, qubits are the smallest unit of [information](#) in [quantum systems](#). Big companies like Google and IBM are competing with research institutes around the world to produce an increasing number of entangled qubits and develop a functioning quantum computer. But a research group at the University of Vienna and the Austrian Academy of Sciences is pursuing a new path to increase the information capacity of complex quantum systems.

The idea behind it is simple: Instead of just increasing the number of particles involved, the complexity of each [system](#) is increased. "The special thing about our experiment is that for the first time, it entangles three photons beyond the conventional two-dimensional nature," explains Manuel Erhard, first author of the study. For this purpose, the Viennese physicists used quantum systems with more than two possible states—in this particular case, the angular momentum of individual light particles. These individual photons now have a higher [information capacity](#) than qubits. However, the entanglement of these light particles turned out to be difficult on a conceptual level. The researchers overcame this challenge with a groundbreaking idea: a computer algorithm that autonomously searches for an experimental implementation.

With the help of a [computer](#) algorithm called Melvin, the researchers found an experimental setup to produce this type of entanglement. At first, this was very complex, but it worked in principle. After some

simplifications, the physicists still faced major technological challenges. The team was able to solve these with state-of-the-art laser technology and a specially developed multi-port. "This multi-port is the heart of our experiment, and combines the three photons so that they are entangled in three dimensions," explains Manuel Erhard.

The peculiar property of the three-photon entanglement in three dimensions allows for experimental investigation of new fundamental questions about the behaviour of quantum systems. In addition, the results of this work could also have a significant impact on future technologies, such as quantum teleportation. "I think the methods and technologies that we developed in this publication allow us to teleport a higher proportion of the total quantum information of a single photon, which could be important for [quantum](#) communication networks," Anton Zeilinger says.

More information: Manuel Erhard et al, Experimental Greenberger–Horne–Zeilinger entanglement beyond qubits, *Nature Photonics* (2018). [DOI: 10.1038/s41566-018-0257-6](https://doi.org/10.1038/s41566-018-0257-6)

Provided by University of Vienna

Citation: One step closer to complex quantum teleportation (2018, November 5) retrieved 23 April 2024 from <https://phys.org/news/2018-11-closer-complex-quantum-teleportation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.