

Breeding corn for water-use efficiency may have just gotten easier

November 14 2018



Water-use efficiency measurements traditionally require specialized equipment, shown here, and are time consuming. Tony Studer's method speeds up the process significantly. Credit: Tony Studer, University of Illinois.

With approximately 80 percent of our nation's water supply going towards agriculture, it's fair to say it takes a lot of water to grow crops. In a climate with less predictable rainfall patterns and more intense droughts, scientists at the University of Illinois are working to reduce water consumption by developing more efficient crops.

"There's a study from many decades ago that shows the amount of water transpired and lost to the air in an acre of [corn](#) is 3 to 4 thousand gallons per day. At 90 million acres of corn in the U.S., plus the length of the growing season, that's lots and lots of water. So there are a lot of improvements that need to be made," says Tony Studer, assistant professor in the Department of Crop Sciences at U of I, and author of a new study in *The Plant Journal*.

A [previous study](#) from Studer's group suggests corn could become 10 to 20 percent more efficient through breeding improvements, which would mean that [plants](#) would be less stressed during short-term droughts. Theoretically, this could add protection for farmers, given uncertain weather patterns. But to make that a reality, according to Studer, the breeding process itself needs to become more efficient.

When attempting to improve a certain trait, in this case water-use efficiency, breeders grow a diverse set of corn lines and screen them to find natural variation in the trait. Once they identify promising individuals, breeders then try to locate key genes that will amplify the trait or integrate the trait into lines of corn with additional desirable qualities.

"It takes a lot of time, space, and effort to produce a productive hybrid," Studer says.

Water-use efficiency is typically measured with an instrument that clamps to leaves and monitors the flux of [carbon dioxide](#) and water vapor moving into and out of the leaf. This process is time-consuming and expensive at large scales, as each measurement can take over an hour.

"If you're going to study water use in a breeding environment or in a field at scale, you need something faster," Studer says.

In their current study, Studer and his colleagues developed a new method to screen hundreds or even thousands of plants without the need for time-consuming field measurements. The method, which tests leaf samples in the lab, takes advantage of the fact that the carbon in carbon dioxide exists in two forms in the atmosphere: a more-abundant and lighter form, ^{12}C ; and a less-abundant and heavier form, ^{13}C .

Once carbon dioxide enters plant leaves, the carbon is incorporated into sugars and plant tissues. Scientists can then measure how much ^{13}C was incorporated compared to ^{12}C . For many plants, the ratio of ^{12}C -to- ^{13}C is indicative of their water-use efficiency. But until now, scientists didn't know if the ratio could reliably reflect water status in corn. Studer's study shows it can.

"We found significant variation in the ^{12}C -to- ^{13}C ratio across 36 diverse lines of corn, and the ^{12}C -to- ^{13}C signature is heritable across environments," he says. "Proving that a trait is inherited and expressed across environments allows a plant breeder to select for this trait and is essential when developing new lines."

The finding, derived from controlled greenhouse trials as well as three

field seasons, provides the efficient method Studer was looking for. And it shows that inbred lines whose carbon ratios are within a certain range may have greater water-use efficiency, although it's too early to say how this will play out in hybrids. Right now, it's enough that the trait appears to be heritable—that alone will be a great help to breeders. But Studer has plans for next steps.

"In a past study, we found there's room for improvement in corn's [water](#)-use efficiency. Here, we're showing that the [trait](#) is measurable and heritable, and we can actually use it to try to make improvements," he says. "The next step is identifying the genes in these regions of the genome that we can manipulate. We've moved all the way from a basic idea of developing the science behind these traits to the point where we can actually make improvements."

More information: Robert J. Twohey et al, Leaf Stable Carbon Isotope Composition Reflects Transpiration Efficiency in *Zea mays*, *The Plant Journal* (2018). [DOI: 10.1111/tpj.14135](https://doi.org/10.1111/tpj.14135)

Provided by University of Illinois at Urbana-Champaign

Citation: Breeding corn for water-use efficiency may have just gotten easier (2018, November 14) retrieved 4 June 2024 from <https://phys.org/news/2018-11-corn-water-use-efficiency-easier.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--