


Devonian integrative stratigraphy and timescale of China

November 16 2018



Representative fossil groups from Devonian. Credit: Science China Press

Studies on the Devonian of China have lasted for about 170 years, and important progress has been made recently. Based on well-studied bio- and chronostratigraphy of the Devonian in South China and adjacent areas, in combination with recent achievements in carbon isotope stratigraphy, event stratigraphy and radioactive isotope ages, researchers

from Nanjing briefly summarize the research history and current status of Devonian chronostratigraphy of China, and for the first time introduce a Devonian integrative stratigraphy framework of China.

The Devonian (419.2-358.9 Ma) is the first geological period of the late Paleozoic, spanning about 60.3 Ma from the end of the Silurian, and its beginning is marked by the First Appearance Datum (FAD) of graptolite *Uncinatograptus uniformis*. During this [time interval](#), the Earth's climate system underwent severe perturbations, characterized by drastic dropdown of atmospheric CO₂ concentration, and gradual transition from the Silurian greenhouse Earth to the Permo-Carboniferous icehouse Earth. The complex pattern of Devonian climate change was controlled by multiple factors at different time scales, and induced 25 global events that are characterized by sea-level rise and fall, ocean anoxic/hypoxic events, and/or biological extinctions/turnovers. Determination of high-resolution Devonian integrative stratigraphy and timescale framework is essential for studies on the complex ocean-land-atmosphere interactions during the Devonian at different temporal and spatial scales.

A review paper by Qie et al. (2018) was published recently in *Science China: Earth Sciences*. Based on well-studied bio- and chronostratigraphy of Devonian in South China and adjacent areas, in combination with recent achievements in carbon isotope stratigraphy, event stratigraphy and radioactive isotope ages, this paper summarizes the research history and current status of Devonian chronostratigraphy of China, and for the first time introduces Devonian integrative stratigraphy framework of China.

In 1985, the global Devonian chronostratigraphy scale was formally defined and subdivided into three series, the lower, middle and upper, and contains seven global stages. Until 1996, all seven Devonian stages' GSSPs have been formally defined and ratified. Conodont is a key taxon

in the Devonian biostratigraphic research. Six out of the seven global stages of the Devonian, except for the basal Devonian boundary, are marked by the FAD of certain conodont species.


By summarizing the taxonomy and biostratigraphy of conodonts in China, Wang (2018) confirmed 58 Devonian conodonts zones in pelagic facies, among which the biozones of Emsian to Famennian can be precisely correlated to the standard international conodont zones. The bases of the Emsian and the Devonian-Carboniferous boundary (DCB) has been defined by the FAD of conodont *Polygnathus kitabicus* and *Siphonodella sulcata*. However, the acceptance of the GSSPs for the bases of the Emsian and DCB have received major criticism since the very beginning of their ratification, and in 2008, the Subcommittee on Devonian Stratigraphy (SDS) and Subcommittee on Carboniferous Stratigraphy (SCCS) have decided to redefine the boundary criteria and levels, which are still ongoing investigations. In addition to conodont biostratigraphy, important progresses on biostratigraphic research of graptolite, ostracods, brachiopods, rugose corals, as well as spores and plants are emphatically introduced in this paper.

Age (Ma)	Global stages		International conodont zone	PC	Boundary age (Ma)	CRS		Biozonation in pelagic facies						
	Se.	Stage				Se.	Stage	Conodont	Ammonoid	Graptolite	Tentaculite	Ostracod		
360	Upper	Famennian	<i>S. sulcata</i>	358.9±0.4 New DCB definition (in discussion)	371.93- 371.78	Upper	Tour.	<i>S. sulcata</i>	<i>Gattendorfia</i>	H. u. ultimus	<i>R. lator</i> <i>R. lator-hemisphaerica</i> int. <i>Maternella hemisphaerica</i> <i>M. dichotoma</i> <i>R. eocostata</i> <i>Richteria serratostrata</i> - <i>Nehdentomis nehdensis</i>			
362			Shandongian				<i>S. praesulcata</i>	<i>C. eurymphala</i>						
364			Yangshuoan				<i>Pa. expansa</i>	<i>Wocklumeria</i>						
366							<i>Pa. postera</i>	<i>Clymenia</i>						
368							<i>Pa. trachytera</i>	<i>Platyclymenia</i>						
370		Frasnian	<i>Pa. marginifera</i>			371.93- 371.78	Upper	Xikuangshanian	<i>Pa. marginifera</i>			<i>Cheiloceras</i>		
372								<i>Pa. rhomboidea</i>						
374								<i>Pa. crepida</i>						
376								<i>Pa. triangularis</i>						
378								<i>Pa. linguiformis</i>	<i>Crickites</i>					
380	Middle	Givetian	<i>Pa. rhenana</i>	382.7±1.6	Middle	Dongganglingian	<i>Pa. rhenana</i>	<i>Archoceras</i>	Homoctenus t. tenuicinctus	<i>En. splendens</i> <i>splen.-R. reichii</i> int. <i>Rabienella reichii</i> <i>Ra. schmidti</i> <i>Rabienella volki</i> <i>Rabienella materni</i> <i>materni/ barrandei</i> <i>bar. riccati</i> int.				
382							<i>Pa. jamieae</i>	<i>Playfordites</i>						
384							<i>Pa. hassi</i>	<i>Beloceras</i>						
386							<i>Pa. punctata</i>	<i>Mesobeloceras</i>						
388							<i>Pa. transitans</i>	<i>Prachorites</i> <i>Proheroceras</i>						
390		Eifelian	<i>Mesotaxis guanyushanensis (=falsiovalis)</i>			387.7±0.8	Middle	Yingtangian			<i>Mesotaxis guanyushanensis (=falsiovalis)</i>	<i>Timanites</i>	<i>N. regularis</i>	<i>B. cicatricosa</i> <i>B. cicatricosa-F. torleyi</i> int.
392											<i>K. disparilis</i>	<i>Koenenites</i>		
394											<i>S. hermanni</i>	<i>Ponticeras</i>		
396											<i>P. varcus</i>	<i>Pharciceras</i>		
398											<i>P. hemiansatus pensensis</i>	<i>M. terebratum</i> <i>Maenioceras molarium</i>		
400	Lower	Emsian	<i>T.k. kockelianus</i>	393.3±1.2	Lower	Sipaian	<i>T.k. kockelianus</i>	<i>Cabrieroceras crispiforme</i>	<i>V. multicostata</i> <i>V. minuta</i> <i>Nowakia (N.) otomari</i> <i>R. nayiensis</i> <i>V. guangxiensis</i> <i>N. albertii</i> <i>N. s. sulcata</i> <i>N. s. procera</i> <i>antiqua</i> <i>holyocera</i> <i>N. muriei</i> <i>N. multicostata</i> <i>N. holynensis?</i>	<i>B. subrecta</i> <i>Bertilionella praeerecta</i> <i>R. longisulcata</i> <i>Bisulco. tuberculata</i>				
402							<i>P. c. costatus</i>	<i>Pinacites-Foordites</i>						
404							<i>P. c. partitus</i>	<i>Anarcestes</i>						
406							<i>P. c. patulus</i>	<i>Anetoceras/Anarcestes</i>						
408			Pragian				<i>P. serotinus</i>	<i>A. (Teneroceras)</i>						
410						<i>P. inversus</i>	<i>A. (Anetoceras)</i>							
412						<i>Polygnathus nothoperbonus</i>	<i>Nandanoceras</i>							
414		Lochkovian				<i>P. gronbergi</i>								
416							<i>P. excavatus</i>							
418						<i>P. kitabicus?</i>								
420			<i>P. pireneae</i>											
422			<i>G. kindlei</i>											
424		<i>G. irregularis</i>												
426		<i>P. gilberti</i>												
428		<i>M. pandora</i>												
430		<i>A. trigonicus</i>												
432		<i>L. transitans</i>												
434		<i>L. elegans</i>												
436		<i>L. omoalpha</i>												
438		<i>Caudicriodus postwoschmidti</i>												
440		<i>Caudicriodus hesperius</i>												
442		<i>D. detorta</i>												

Devonian Chronostratigraphy and biozonations in pelagic facies in China.
Credit: Science China Press

As for chemostratigraphy, carbon and strontium isotopic records from South China show a similar trend with Euramerica sections, and have global correlation significances. $\delta^{13}\text{C}$ values show major positive shifts in the Silurian-Devonian boundary, Kačák, Frasnian, lower and upper Kellwasser, as well as Hangenberg events, suggesting global carbon cycle perturbation during these critical time intervals. In Longmenshan region, $^{87}\text{Sr}/^{86}\text{Sr}$ values are 0.70788-0.70868, and its long-term trend agrees with a published mean LOWESS fitted line. After a gradual decline from early Devonian, $^{87}\text{Sr}/^{86}\text{Sr}$ ratios reach nadir values and stay stable in middle Devonian, then start to increase in Frasnian and define a plateau throughout the Famennian. $^{87}\text{Sr}/^{86}\text{Sr}$ ratios can be used as a tool for precise stratigraphic correlation and dating. This paper also gives a detail introduction to Devonian event stratigraphy, cyclostratigraphy, geological chronology, and stratigraphic correlation among main major palaeobiogeography realms in China.

At present, the main tasks for researchers of the Devonian of China include: (1) establishment of continuous astronomical time scale and high-resolution radioactive isotope dating; (2) further subdivision of Chinese regional chronostratigraphy; (3) sulfur and oxygen isotopic stratigraphy of Devonian; (4) integrated stratigraphy studies on the Devonian System in other stratigraphic regions except South China region.



Generalized Devonian carbon and strontium isotope stratigraphy, event stratigraphy in China. Credit: Science China Press

More information: Wenkun Qie et al, Devonian integrative stratigraphy and timescale of China, *Science China Earth Sciences* (2018).
[DOI: 10.1007/s11430-017-9259-9](https://doi.org/10.1007/s11430-017-9259-9)

Provided by Science China Press

Citation: Devonian integrative stratigraphy and timescale of China (2018, November 16)
retrieved 25 April 2024 from
<https://phys.org/news/2018-11-devonian-stratigraphy-timescale-china.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.