

eDNA emerges as powerful tool for tracking threatened river herring in Chesapeake Bay

November 1 2018

Using environmental DNA (eDNA) to track the presence of fish in waterways is emerging as a powerful tool to detect and understand the abundance of species in aquatic environments. However, relatively few studies have compared the performance of this emerging technology to traditional catch or survey approaches in the field.

Researchers from the University of Maryland Center for Environmental Science and Smithsonian Environmental Research Center field tested using eDNA—tracking the presence of fish by identifying DNA that has been left behind in the water—to detect [river herring](#) in tributaries of Maryland's Chesapeake Bay. They found that tracking and quantifying [herring](#) DNA from the environment corresponded well to more traditional field methods and has great potential to assist future monitoring efforts of river herring abundance and habitat use.

"Sampling a single river, you need a net, crew, permits, it can be expensive," said study author Louis Plough of the University of Maryland Center for Environmental Science. "The eDNA approach is an alternative where you just take water and you get an idea of the abundance of fish."

Accurate information on how many fish are in a river system and where they travel is essential to understanding their ecology and how to manage the fishery in increasingly human-impacted environments.

Environmental DNA (eDNA) sampling has recently emerged as a powerful, non-invasive alternative to capture-based techniques to detect

the presence of a species in their environment.

"You can scoop up water and know what's been there," said Plough. "It captures a snapshot of the DNA that has been around in the past couple of weeks."

The researchers developed a genetic probe that targeted and identified the DNA of two particular fish species, alewife and blueback herring, collectively known as river herring, in the waterways. As they swim, the fish slough off cells and waste that contain DNA.

"The eDNA method is especially powerful because it can identify whether alewife, blueback herring, or both species are present at each site," said study co-author Matt Ogburn of the Smithsonian Environmental Research Center. "When we use some traditional methods like collecting fish eggs or larvae, we often aren't able to tell which river herring species they belong to, and they might even be from other local species like hickory shad."

River herring, the common name for alewife and blueback herring, were once common in Maryland's rivers, where they came to spawn every spring. One of the oldest fisheries in North America, they play important roles in coastal food webs—striped bass, bald eagles and osprey like to snack on them—yet the species has declined sharply since the 1970s due to loss of spawning habitat, overfishing, and degradation of water quality.

"These species live in saltwater and spawn in freshwater every spring. They've been hit hard by overfishing and habitat loss, including dams that blocked their way to spawning grounds upstream, so numbers have declined," said Plough. "We need to conserve them, they need help, but it's expensive to monitor them."

The team sampled 196 locations across 12 tributaries in Maryland—nearly 500 waters samples—in an effort to understand where they are and how many there are.

They found that the eDNA abundance data corresponded well to other field methods and even gave them a finer looks at species distribution. For example, alewife were more common among eastern shore rivers while blueback herring were more common among rivers on the more developed western shore.

The researchers are among first to use eDNA across the Chesapeake Bay watershed to create a map of where river herring are spawning.

"The only other large-scale surveys of river herring spawning habitats in the Chesapeake Bay watershed were conducted in Maryland and Virginia from the 1960s to 1980s, and different methods were used in each state," Ogburn said. "Our study is the first to use the same method across the watershed, which allows us to study habitat preferences at an unprecedented scale."

"The goal is to do more of this with other species, using new tools to inform conservation and restoration efforts," said Plough.

More information: Louis V. Plough et al. Environmental DNA analysis of river herring in Chesapeake Bay: A powerful tool for monitoring threatened keystone species, *PLOS ONE* (2018). [DOI: 10.1371/journal.pone.0205578](https://doi.org/10.1371/journal.pone.0205578)

Provided by University of Maryland Center for Environmental Science

Citation: eDNA emerges as powerful tool for tracking threatened river herring in Chesapeake

Bay (2018, November 1) retrieved 18 April 2024 from <https://phys.org/news/2018-11-edna-emerges-powerful-tool-tracking.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.