

# Graphene on the way to superconductivity

#### November 9 2018



The bandstructure in a double layer of graphene was scanned by ARPES at synchrotron light source BESSY II. Credit: HZB



Scientists at HZB have found evidence that double layers of graphene have a property that may let them conduct current completely without resistance. They probed the band structure at BESSY II with extremely high resolution ARPES and could identify a flat area at a surprising location. Their research is published in *Science Advances*.

Carbon atoms can form bonds in multiple ways. Pure carbon can therefore occur in many forms, including diamond, graphite, nanotubes, football molecules or as a honeycomb net with hexagonal meshes, known as <u>graphene</u>. This exotic, strictly two-dimensional material conducts electricity well, but is not a superconductor. But perhaps this can be changed.

## A complicated option for superconductivity

In April 2018, a group at MIT in the U.S. showed that it is possible to generate a form of superconductivity in a system of two layers of graphene under very specific conditions. To do this, the two hexagonal nets must be twisted against each other at a 1.1 degree angle. Under this condition, a flat <u>band</u> forms in the electronic structure. The preparation of samples from two layers of graphene with such an exactly adjusted twist is complex, and not suitable for mass production. Nevertheless, the study has attracted a lot of attention among experts.

### The simple way to flat bands

But there is one more, much simpler way of flat band formation. This was shown by a group at the HZB around Prof. Oliver Rader and Dr. Andrei Varykhalov with investigations at BESSY II.

The samples were provided by Prof. Thomas Seyller, TU Chemnitz. There they are produced using a process that is also suitable for the



production of larger areas and in large quantities: A silicon carbide crystal is heated until silicon atoms evaporate from the surface, leaving first a single-layer of graphene on the surface, and then a second layer of graphene. The two graphene layers are not twisted against each other, but lie exactly on top of each other.

#### Scanning the band structure with ARPES

At BESSY II, the physicists are able to scan the so-called band structure of the sample. This band structure provides information on how the charge carriers are distributed among the quantum-mechanically permitted states and which charge carriers are available for transport at all. The angle-resolved photoemission spectroscopy (ARPES) at BESSY II enables such measurements with extremely high resolution.

Via an exact analysis of the band structure, they identified an area that had previously been overlooked. "The double layer of graphene has been studied before because it is a semiconductor with a band gap," explains Varykhalov. "But on the ARPES instrument at BESSY II, the resolution is high enough to recognize the flat area next to this band gap."

"It is an overseen property of a well-studied system," says first author Dr. Dmitry Marchenko. "It was previously unknown that there is a flat area in the band <u>structure</u> in such a simple well-known system."

This flat area is a prerequisite for superconductivity, but only if it is situated exactly at the so-called Fermi energy. In the case of the two-<u>layer</u> graphene, its energy level is only 200 milli-electron volts below the Fermi energy, but it is possible to raise the energy level of the flat area to the Fermi energy either by doping with foreign atoms or by applying an external voltage, the so-called gate voltage.

The physicists have found that the interactions between the two graphene



<u>layers</u> and between graphene and the silicon carbide lattice are jointly responsible for the formation of the flat band area. "We can predict this behavior with very few parameters and could use this mechanism to control the <u>band structure</u>," adds Oliver Rader.

**More information:** "Extremely flat band in bilayer graphene" *Science Advances* (2018). DOI: 10.1126/sciadv.aau0059, advances.sciencemag.org/content/4/11/eaau0059

#### Provided by Helmholtz Association of German Research Centres

Citation: Graphene on the way to superconductivity (2018, November 9) retrieved 10 May 2024 from <u>https://phys.org/news/2018-11-graphene-superconductivity.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.