

# Hubble uncovers thousands of globular star clusters scattered among galaxies

November 29 2018

---



This is a Hubble Space Telescope mosaic of the immense Coma cluster of over 1,000 galaxies, located 300 million light-years from Earth. Hubble's incredible sharpness was used to do a comprehensive census of the cluster's most diminutive members: a whopping 22,426 globular star clusters (circled in green). Among the earliest homesteaders of the universe, globular star clusters are snow-globe-shaped islands of several hundred thousand ancient stars. The survey found the globular clusters scattered in the space between the galaxies. They have been orphaned from their home galaxies through galaxy tidal interactions within the bustling cluster. Astronomers will use the globular cluster field for mapping the distribution of matter and dark matter in the Coma galaxy cluster. Credit:

NASA, ESA, J. Mack (STScI), and J. Madrid (Australian Telescope National Facility)

Gazing across 300 million light-years into a monstrous city of galaxies, astronomers have used NASA's Hubble Space Telescope to do a comprehensive census of some of its most diminutive members: a whopping 22,426 globular star clusters found to date.

The survey, published in the November 9, 2018, issue of *The Astrophysical Journal*, will allow for astronomers to use the globular cluster field to map the distribution of matter and dark matter in the Coma galaxy cluster, which holds over 1,000 galaxies that are packed together.

Because [globular clusters](#) are much smaller than entire galaxies – and much more abundant – they are a much better tracer of how the fabric of space is distorted by the Coma cluster's gravity. In fact, the Coma cluster is one of the first places where observed gravitational anomalies were considered to be indicative of a lot of unseen mass in the universe – later to be called "dark matter."

Among the earliest homesteaders of the universe, globular star clusters are snow-globe-shaped islands of several hundred thousand ancient stars. They are integral to the birth and growth of a galaxy. About 150 globular clusters zip around our Milky Way galaxy, and, because they contain the oldest known stars in the universe, were present in the early formative years of our galaxy.

Some of the Milky Way's globular clusters are visible to the [naked eye](#) as fuzzy-looking "stars." But at the distance of the Coma cluster, its globulars appear as dots of light even to Hubble's super-sharp vision. The

survey found the globular clusters scattered in the space between the galaxies. They have been orphaned from their home galaxy due to galaxy near-collisions inside the traffic-jammed cluster. Hubble revealed that some globular clusters line up along bridge-like patterns. This is telltale evidence for interactions between galaxies where they gravitationally tug on each other like pulling taffy.

Astronomer Juan Madrid of the Australian Telescope National Facility in Sydney, Australia first thought about the distribution of globular clusters in Coma when he was examining Hubble images that show the globular clusters extending all the way to the edge of any given photograph of galaxies in the Coma cluster.

He was looking forward to more data from one of the legacy surveys of Hubble that was designed to obtain data of the entire Coma cluster, called the Coma Cluster Treasury Survey. However, halfway through the program, in 2006, Hubble's powerful Advanced Camera for Surveys (ACS) had an electronics failure. (The ACS was later repaired by astronauts during a 2009 Hubble servicing mission.)

To fill in the [survey](#) gaps, Madrid and his team painstakingly pulled numerous Hubble images of the galaxy cluster taken from different Hubble observing programs. These are stored in the Space Telescope Science Institute's Mikulski Archive for Space Telescopes in Baltimore, Maryland. He assembled a mosaic of the central region of the cluster, working with students from the National Science Foundation's Research Experience for Undergraduates program. "This program gives an opportunity to students enrolled in universities with little or no astronomy to gain experience in the field," Madrid said.

The team developed algorithms to sift through the Coma mosaic images that contain at least 100,000 potential sources. The program used globular clusters' color (dominated by the glow of aging red stars) and

spherical shape to eliminate extraneous objects – mostly background [galaxies](#) unassociated with the Coma [cluster](#).

Though Hubble has superb detectors with unmatched sensitivity and resolution, their main drawback is that they have tiny fields of view.

"One of the cool aspects of our research is that it showcases the amazing science that will be possible with NASA's planned Wide Field Infrared Survey Telescope (WFIRST) that will have a much larger field of view than Hubble," said Madrid. "We will be able to image entire galaxy clusters at once."

**More information:** Juan P. Madrid et al. A Wide-field Map of Intracluster Globular Clusters in Coma, *The Astrophysical Journal* (2018). [DOI: 10.3847/1538-4357/aae206](https://doi.org/10.3847/1538-4357/aae206)

Provided by ESA/Hubble Information Centre

Citation: Hubble uncovers thousands of globular star clusters scattered among galaxies (2018, November 29) retrieved 18 April 2024 from <https://phys.org/news/2018-11-hubble-uncovers-thousands-globular-star.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.