

Human ancestors not to blame for ancient mammal extinctions in Africa

November 22 2018



Africa contains the greatest diversity of large-bodied mammals in the world today. This includes five species of megaherbivores (herbivores over 2000 lbs). From left to right: hippopotamus (*Hippopotamus amphibius*), giraffe (*Giraffa camelopardalis*), elephant (*Loxodonta africana*), white rhino (*Ceratotherium simum*), and black rhino (*Diceros bicornis*). Credit: John Rowan

New research disputes a long-held view that our earliest tool-bearing ancestors contributed to the demise of large mammals in Africa over the last several million years. Instead, the researchers argue that long-term environmental change drove the extinctions, mainly in the form of grassland expansion likely caused by falling atmospheric carbon dioxide (CO₂) levels.

Tyler Faith, curator of archaeology at the Natural History Museum of

Utah and assistant professor in the Department of Anthropology at the University of Utah, led the study. The research team also includes John Rowan from the University of Massachusetts Amherst, Andrew Du from the University of Chicago, and Paul Koch from the University of California, Santa Cruz.

The study is published today in the journal *Science*.

"Despite decades of literature asserting that early hominins impacted ancient African faunas, there have been few attempts to actually test this scenario or to explore alternatives," Faith says. "We think our study is a major step towards understanding the depth of anthropogenic impacts on large mammal communities, and provides a convincing counter-argument to these long-held views about our early ancestors."

To test for ancient hominin impacts, the researchers compiled a seven-million-year record of herbivore extinctions in eastern Africa, focusing on the very largest [species](#), the so-called 'megaherbivores' (species over 2,000 lbs.) Though only five megaherbivores exist in Africa today, there was a much greater diversity in the past. For example, three-million-year-old 'Lucy' (*Australopithecus afarensis*) shared her woodland landscape with three giraffes, two rhinos, a hippo, and four elephant-like species at Hadar, Ethiopia.

When and why these species disappeared has long been a mystery for archaeologists and paleontologists, despite the evolution of tool-using and meat-eating hominins getting most of the blame.

"Our analyses show that there is a steady, long-term decline of megaherbivore diversity beginning around 4.6 million years ago. This extinction process kicks in over a million years before the very earliest evidence for human ancestors making tools or butchering animal carcasses and well before the appearance of any hominin species

realistically capable of hunting them, like *Homo erectus*," says Faith.




The team analyzed more than 100 sites in East Africa with rich fossil records to track the longterm decline of megaherbivore diversity. Credit: J. Tyler Faith

Taking a Closer Look

Faith and his team quantified long-term changes in eastern African megaherbivores using a dataset of more than 100 fossil assemblages spanning the last seven million years. The team also examined independent records of climatic and environmental trends and their effects, specifically global atmospheric CO₂, stable carbon isotope records of vegetation structure, and stable carbon isotopes of eastern African fossil herbivore teeth, among others.

Their analysis reveals that over the last seven million years substantial megaherbivore extinctions occurred: 28 lineages became extinct, leading to the present-day communities lacking in large animals. These results highlight the great diversity of ancient megaherbivore communities, with many having far more megaherbivore species than exist today across Africa as a whole.




A fossil tooth of a hippo (*Hippopotamus amphibius*) (left) and a fossil tooth of a white rhinoceros (*Ceratotherium simum*) (right), two of the few surviving

megaherbivores, from the Late Pleistocene of western Kenya (left). Credit: J. Tyler Faith

Further analysis showed that the onset of the megaherbivore decline began roughly 4.6 million years ago, and that the rate of diversity decline did not change following the appearance of *Homo erectus*, a human ancestor often blamed for the extinctions. Rather, Faith's team argues that climate is more likely culprit.

"The key factor in the Plio-Pleistocene megaherbivore decline seems to be the expansion of grasslands, which is likely related to a global drop in atmospheric CO₂ over the last five million years," says John Rowan, a postdoctoral scientist from University of Massachusetts Amherst. "Low CO₂ levels favor tropical grasses over trees, and as a consequence savannas became less woody and more open through time. We know that many of the extinct megaherbivores fed on woody vegetation, so they seem to disappear alongside their food source."

The loss of massive herbivores may also account for other extinctions that have also been attributed to ancient hominins. Some scientist suggest that competition with increasingly carnivorous species of *Homo* led to the demise of numerous carnivores over the last few million years. Faith and his team suggest an alternative.



The decline of African megaherbivore diversity (gray curve) over the last seven million years was driven by falling atmospheric carbon dioxide and the expansion of grasslands, not ancient hominin impacts. The onset of the megaherbivore decline around 4.6 million years ago (red dashed line and shading) occurs well before the appearance of tool-bearing hominin species capable of hunting large prey. Credit: John Rowan

"We know there are also major extinctions among African carnivores at this time and that some of them, like saber-tooth cats, may have specialized on very large prey, perhaps juvenile elephants" says Paul Koch. "It could be that some of these carnivores disappeared with their megaherbivore prey."

"Looking at all of the potential drivers of the megaherbivore decline, our analyses suggest that changing climate and environment played the key role in Africa's past extinctions," said Faith. "It follows that in the search for ancient hominin impacts on ancient African ecosystems, we must focus our attention on the one species known to be capable of causing them—us, *Homo sapiens*, over the last 300,000 years."

More information: J.T. Faith at University of Utah in Salt Lake City, UT et al., "Plio-Pleistocene decline of African megaherbivores: No evidence for ancient hominin impacts," *Science* (2018).

[science.sciencemag.org/cgi/doi ... 1126/science.aau2728](https://science.sciencemag.org/cgi/doi/10.1126/science.aau2728)

R. Bobe et al., "The decline of Africa's largest mammals," *Science* (2018). [science.sciencemag.org/cgi/doi ... 1126/science.aav6883](https://science.sciencemag.org/cgi/doi/10.1126/science.aav6883)

Provided by University of Utah

Citation: Human ancestors not to blame for ancient mammal extinctions in Africa (2018, November 22) retrieved 27 April 2024 from <https://phys.org/news/2018-11-human-ancestors-blame-ancient-mammal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--