


What could cause the Mississippi Bight to become hypoxic?

November 8 2018



Mississippi Bight region with the CONCORDE 2016 summer survey transects (labelled thick red lines) with the associated dissolved oxygen (DO) conditions. Coloration is 7-d average of chlorophyll-a derived from MODIS sensor (1 km resolution) for the week of July 20-26, 2016. Due to the optical complexity of coastal waters, the chlorophyll-a concentration represents relative values but effectively illustrates an intrusion of fresher water (yellow filament) on the shelf

in contrast to clear Gulf of Mexico water (blue). Credit: *Continental Shelf Research*

Coastal regions with low dissolved oxygen (known as hypoxia) can lead to poor water quality and harm regional fisheries. These areas of low dissolved oxygen are expanding and expected to continue growing in coming years due to human impacts on the environment.

A recent article published in *Continental Shelf Research* explores aspects of the environmental conditions that can potentially lead to hypoxia in the Mississippi Bight [region](#) of the northern Gulf of Mexico. This area extends from Apalachicola in Florida to the Mississippi River Delta.

Dr. Brian Dzwonkowski of the Dauphin Island Sea Lab and University of South Alabama worked with several other research groups to synthesize a range of data sets. The article, "Tracking sea surface salinity and dissolved [oxygen](#) on a river-influenced, seasonally stratified shelf, Mississippi Bight, northern Gulf of Mexico" was published in September 2018.

Understanding the Dead Zone

The northern Gulf of Mexico is home to the second largest hypoxic zone in the world, also known as the "Dead Zone", however, there is no part of the country or the world that is immune. Researchers have directly linked the northern Gulf of Mexico "Dead Zone" to discharge from the Mississippi River. The river water delivers excess nutrients to the region and creates a fresher layer of water at the surface that prevents oxygen from reaching the bottom layer. This leads to a depletion of oxygen over the summer season.

The historic and current focus on "Dead Zone" research and management is centered on the Louisiana and Texas shelf region of the northern Gulf of Mexico, a region to the west of the Mississippi Delta.

The Research

Using satellite [sea surface salinity](#) data from 2010 to 2016, Dr. Dzwonkowski and his colleagues demonstrated the region to the east of the delta is similarly impacted by river discharge. Accompanying field data from the July of 2016, showed evidence of widespread areas of low bottom dissolved oxygen (and hypoxic conditions in some areas) across the Mississippi Bight region.

As a result, there are potentially large regions of hypoxic and low dissolved oxygen zones that are not included in the yearly estimate of the "Dead Zone".

Furthermore, the low dissolved oxygen levels observed, as well as the extensive freshwater surface layer, suggest this region is highly susceptible to becoming hypoxic should there be changes to background environmental conditions (e.g., increased ocean warming), regional watershed land use (e.g., coastal urbanization), and/or diversion pathways of the Mississippi River (e.g., increased discharge through spillways).

More information: Brian Dzwonkowski et al, Tracking sea surface salinity and dissolved oxygen on a river-influenced, seasonally stratified shelf, Mississippi Bight, northern Gulf of Mexico, *Continental Shelf Research* (2018). [DOI: 10.1016/j.csr.2018.09.009](https://doi.org/10.1016/j.csr.2018.09.009)

Provided by Dauphin Island Sea Lab

Citation: What could cause the Mississippi Bight to become hypoxic? (2018, November 8)
retrieved 23 April 2024 from <https://phys.org/news/2018-11-mississippi-bight-hypoxic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.