

NIST atomic clocks now keep time well enough to improve models of Earth

November 28 2018



NIST physicist Andrew Ludlow and colleagues achieved new atomic clock performance records in a comparison of two ytterbium optical lattice clocks. Laser systems used in both clocks are visible in the foreground, and the main apparatus for one of the clocks is located behind Ludlow. Credit: Burrus/NIST

Experimental atomic clocks at the National Institute of Standards and Technology (NIST) have achieved three new performance records, now



ticking precisely enough to not only improve timekeeping and navigation, but also detect faint signals from gravity, the early universe and perhaps even dark matter.

The clocks each trap a thousand <u>ytterbium atoms</u> in optical lattices, grids made of laser beams. The atoms tick by vibrating or switching between two <u>energy levels</u>. By comparing two independent clocks, NIST physicists achieved record performance in three important measures: systematic uncertainty, stability and reproducibility.

Published online Nov. 28 in the journal *Nature*, the new NIST <u>clock</u> records are:

- Systematic uncertainty: How well the clock represents natural vibrations, or <u>frequency</u>, of the atoms. NIST researchers found that each clock ticked at a rate matching the natural frequency to within a possible error of just 1.4 parts in 10¹⁸—about one billionth of a billionth.
- Reproducibility: How closely the two clocks tick at the same frequency, shown by 10 comparisons of the clock pair, yielding a frequency difference below the 10⁻¹⁸level (again, less than one billionth of a billionth).

"Systematic uncertainty, stability, and reproducibility can be considered the 'royal flush' of performance for these clocks," project leader Andrew Ludlow said. "The agreement of the two clocks at this unprecedented level, which we call reproducibility, is perhaps the single most important result, because it essentially requires and substantiates the other two results."



"This is especially true because the demonstrated reproducibility shows that the clocks' total error drops below our general ability to account for gravity's effect on time here on Earth. Hence, as we envision clocks like these being used around the country or world, their relative performance would be, for the first time, limited by Earth's gravitational effects."

Einstein's theory of relativity predicts that an atomic clock's ticking, that is, the frequency of the atoms' vibrations, is reduced—shifted toward the red end of the electromagnetic spectrum—when operated in stronger gravity. That is, time passes more slowly at lower elevations.

While these so-called redshifts degrade a clock's timekeeping, this same sensitivity can be turned on its head to exquisitely measure gravity. Super-sensitive clocks can map the gravitational distortion of space-time more precisely than ever. Applications include relativistic geodesy, which measures the Earth's gravitational shape, and detecting signals from the early universe such as gravitational waves and perhaps even asyet-unexplained "dark matter."

NIST's ytterbium clocks now exceed the conventional capability to measure the geoid, or the shape of the Earth based on tidal gauge surveys of sea level. Comparisons of such clocks located far apart such as on different continents could resolve geodetic measurements to within 1 centimeter, better than the current state of the art of several centimeters.

In the past decade of new clock performance records announced by NIST and other labs around the world, this latest paper showcases reproducibility at a high level, the researchers say. Furthermore, the comparison of two clocks is the traditional method of evaluating performance.

Among the improvements in NIST's latest ytterbium clocks was the inclusion of thermal and electric shielding, which surround the atoms to



protect them from stray electric fields and enable researchers to better characterize and correct for frequency shifts caused by heat radiation.

The ytterbium atom is among potential candidates for the future redefinition of the second—the international unit of time—in terms of optical frequencies. NIST's new clock records meet one of the international redefinition roadmap's requirements, a 100-fold improvement in validated accuracy over the best clocks based on the current standard, the cesium atom, which vibrates at lower microwave frequencies.

NIST is building a portable ytterbium lattice clock with state-of-the-art performance that could be transported to other labs around the world for clock comparisons and to other locations to explore relativistic geodesy techniques.

More information: W. F. McGrew et al. Atomic clock performance enabling geodesy below the centimetre level, *Nature* (2018). DOI: 10.1038/s41586-018-0738-2

Provided by National Institute of Standards and Technology

Citation: NIST atomic clocks now keep time well enough to improve models of Earth (2018, November 28) retrieved 27 April 2024 from <u>https://phys.org/news/2018-11-nist-atomic-clocks-earth.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.